

NESTING AND BROODING CHARACTERISTICS OF SHARP-TAILED  
GROUSE (PEDIOECETES PHASIANELLUS JAMESI LINCOLN) IN

SOUTHWESTERN NORTH DAKOTA

by

Carter Drew Christenson

Bachelor of Science, University of North Dakota, 1970

A Thesis

Submitted to the Faculty

of the

University of North Dakota

in partial fulfillment of the requirements

for the Degree of

Master of Science

Grand Forks, North Dakota

December

1970

\*12930047  
D88018676

-III-

OL  
696  
.52  
C474  
1970

This thesis submitted by Carter Drew Christenson in partial fulfillment of the requirements for the Degree of Master of Science from the University of North Dakota is hereby approved by the Faculty Advisory Committee under whom the work has been done.

BLM LIBRARY  
SC-324A, BLDG. 50  
DENVER FEDERAL CENTER  
P. O. BOX 25047  
DENVER, CO 80225-0047

James R. Reilly  
(Chairman)

John B. Owen

Dean W. Blinn

A. William Johnson  
Dean of the Graduate School

Permission

Title Nesting and Brooding Characteristics of Sharp-tailed Grouse  
(Pedioecetes phasianellus jamesi Lincoln) in Southwestern North  
Dakota  
Department Biology  
Degree Master of Science

In presenting this thesis in partial fulfillment of the requirements for a graduate degree from the University of North Dakota, I agree that the Library of this University shall make it freely available for inspection. I further agree that permission for extensive copying for scholarly purposes may be granted by the professor who supervised my thesis work or, in his absence by the Chairman of the Department or the Dean of the Graduate School. It is understood that any copying or publication or other use of this thesis or part thereof for financial gain shall not be allowed without my written permission. It is also understood that due recognition shall be given to me and to the University of North Dakota in any scholarly use which may be made of any material in my thesis.

Signature Carter D. Christenson

Date December 10, 1970

ACKNOWLEDGEMENTS

I wish to thank Commissioner Russell W. Stuart and Deputy Commissioner Wilbur Boldt, North Dakota State Game and Fish Department<sup>1</sup>, for the opportunity to conduct this research.

To Dr. James R. Reilly, my Advisor, I express my deepest gratitude for his untiring efforts in guiding the research and inspecting and refining this thesis. I am grateful to Dr. Dean W. Blinn and Dr. John B. Owen for serving on my committee and making valuable suggestions throughout the study.

Gratitude is expressed to Gerald D. Kobriger and C. R. (Chris) Grondahl, North Dakota State Game and Fish Department, for their continual assistance and encouragement throughout the field investigation. Also, special thanks is extended to Bruce Dreher and Dennis Johnson, Medora Ranger District, U. S. Forest Service, for their technical assistance in the analysis and sampling of vegetation at nesting sites.

I wish to thank Dr. Warren Whitman, North Dakota State University, for his valuable assistance in identifying nest site vegetation and making project suggestions. And to my wife, Gwen, for her constant reassurance and continual assistance in both research and manuscript preparation, my most sincere thanks.

---

<sup>1</sup>A contribution in part of Federal Aid in Wildlife Restoration, Pittman-Rortson Project W-67-R-10 and 11, Job 13, North Dakota.

ACK

LIS

LIS

ABS

INT

DES

LIT

MET

RES

DIS

REC

LIT



TABLE OF CONTENTS

|                                     |           |
|-------------------------------------|-----------|
| ACKNOWLEDGEMENTS                    | Page<br>V |
| LIST OF TABLES                      | VII-VIII  |
| LIST OF FIGURES                     | IX        |
| ABSTRACT                            | X         |
| INTRODUCTION                        | I - 1     |
| DESCRIPTION OF THE STUDY AREA       | 1         |
| Location                            |           |
| Native vegetation                   |           |
| Land-use                            |           |
| LITERATURE REVIEW                   | 4-9       |
| Grouse habitat and range management |           |
| Telemetry                           |           |
| Trapping techniques                 |           |
| Hen dispersal                       |           |
| Nesting characteristics             |           |
| Predation                           |           |
| Brood habitat                       |           |
| Vegetation analysis                 |           |
| METHODS AND MATERIALS               | 9-10      |
| Trapping and marking                |           |
| Telemetry                           |           |
| Dispersal of hens                   |           |
| Vegetation at nest sites            |           |
| Brood movement and cover selection  |           |
| RESULTS                             | 10-42     |
| Trapping and marking                |           |
| Telemetry                           |           |
| Dispersal of hens                   |           |
| Vegetation at nest site             |           |
| Brood movement and cover selection  |           |
| DISCUSSION                          | 43-45     |
| Trapping                            |           |
| Telemetry                           |           |
| Hen dispersal and nesting           |           |
| Vegetation at nest sites            |           |
| Predation                           |           |
| Brood movement and cover selection  |           |
| RECOMMENDATIONS                     | 46        |
| LITERATURE CITED                    | 47-53     |

LIST OF TABLES

| Table |   | Page |
|-------|---|------|
| 1.    | Land-use on 23,680 Acres on and Surrounding Gorham Study Area, Summer, 1968 (Bernhoft, 1969).   | 3    |
| 2.    | Male Sharp-tailed Grouse Census, Gorham Study Area, North Dakota, Spring 1966-1970  | 11   |
| 3.    | Weights of Adult and Immature Sharp-tailed Grouse, Gorham Study Area, North Dakota, 1969-1970   | 11   |
| 4.    | Sharp-tailed Grouse Trapping Methods and Success, Gorham Study Area, North Dakota, 1969-1970  | 12   |
| 5.    | Comparison of Initial Spring and Mid-incubation Weights of Six Sharp-tailed Grouse Hens, Gorham Study Area, North Dakota, 1969-1970   | 12   |
| 6.    | Sharp-tailed Grouse Hen Dispersal and Nesting, Gorham Study Area, North Dakota, 1969-1970   | 15   |
| 7.    | Telemetric Study and Final Disposition of 18 Sharp-tailed Grouse Hens, Gorham Study Area, North Dakota, 1969-1970   | 16   |
| 8.    | Species Composition and Amount of Vegetation at Channel 4-69 and 10-69 Nest Sites, Gorham Study Area, North Dakota. Vegetation Sampled by Clipped Hoop Method, Fall 1969 and Spring 1970                                    | 20   |
| 9.    | Species Composition and Amount of Vegetation at Channel 1-69, 3-69, and 7-69 Nest Sites, Gorham Study Area, North Dakota. Vegetation Sampled by Clipped Hoop Method, Spring 1970  | 21   |
| 10.   | Species Composition and Amount of Vegetation at Channel 6-69, 2-69, and 8 <sup>a</sup> -69 Nest Sites, Gorham Study Area, North Dakota. Vegetation Sampled by Clipped Hoop Method, Spring 1970.                             | 22   |
| 11.   | Species Composition and Amount of Vegetation at Channel 1 <sup>a</sup> -70, 1 <sup>b</sup> -70 and 2-70 Nest Sites, Gorham Study Area, North Dakota. Vegetation Sampled by Clipped Hoop Method, Summer 1970                 | 23   |
| 12.   | Species Composition and Amount of Vegetation at Channel 3 <sup>a</sup> -70, 3 <sup>b</sup> -70, and 6-70 Nest Sites, Gorham Study Area, North Dakota. Vegetation Sampled by Clipped Hoop Method, Summer 1970.               | 24   |
| 13.   | Species Composition and Amount of Vegetation at Channel 10 <sup>a</sup> -70, 10(nest bowl)-70, and 10 <sup>b</sup> -70 Nest Sites, Gorham Study Area, North Dakota. Vegetation Sampled by Clipped Hoop Method, Summer 1970. | 25   |

| Table   | Page  |
|---|-------|
| 14. Species Composition and Amount of Vegetation at Channel 8-70, 12 <sup>a</sup> -70, and 10 <sup>1</sup> -70 Nest Sites Gorham Study Area, North Dakota. Vegetation Sampled by Clipped Hoop Method, Summer 1970 | 26    |
| 15. Characteristics of Cover at Sharp-tailed Grouse Nest Sites, Gorham Study Area, North Dakota, Based on 2 X 1 Foot Cover Board, 1969-1970   | 27-29 |
| 16. Cover Characteristics and Adjacent Land-use at 37 Sharp-tailed Grouse Brood Locations, Gorham Study Area, Southwestern North Dakota, 1969-1970  | 31    |
| 17. Home Range and Daily Movements of Eight Sharp-tailed Grouse Broods, Gorham Study Area, North Dakota, 1969-1970  | 41    |

LIST OF FIGURES

| Figure   | Page |
|--|------|
| 1.. Location map of Gorham Study Area, North Dakota, 1969-1970   | 2    |
| 2. Hen Dispersal from Dancing Grounds, Gorham Study Area, North Dakota, 1969-1970                                  | 14   |
| 3. Channel 2-69 Brood Movement, Gorham Study Area, North Dakota, 1969  | 32   |
| 4. Channel 8-69 Brood Movement, Gorham Study Area, North Dakota, 1969.   | 33   |
| 5. Channel 10-69 Brood Movement, Gorham Study Area, North Dakota, 1969.  | 34   |
| 6. Channel 1-70 Brood Movement, Gorham Study Area, North Dakota, 1970  | 35   |
| 7. Channel 2-70 Brood Movement, Gorham Study Area, North Dakota, 1970  | 37   |
| 8. Channel 6-70 Brood Movement, Gorham Study Area, North Dakota, 1970  | 38   |
| 9. Channel 8-70 Brood Movement, Gorham Study Area, North Dakota, 1970  | 39   |
| 10. Channel 10 <sup>i</sup> -70 Brood Movement, Gorham Study Area, North Dakota, 1970                              | 40   |
| 11. Daily Activities and Cover Selection of Sharp-tailed Grouse Broods, Gorham Study Area, North Dakota, 1969-1970 | 42   |

A B S T R A C T

A study of the nesting and brood habitat preferences of sharp-tailed grouse (Pedioecetes phasianellus jamesi Lincoln) was conducted in southwestern North Dakota during the reproductive seasons of 1969 and 1970. Radio telemetry was used to monitor the spring and summer activities of seventeen sharptail hens. Eight brooding hens were semi-continuously radio-tracked for movement and cover selection. A quantitative and qualitative analysis of seventeen nests and eight renests was made to determine the preferred nesting cover and the effects of intensive grazing upon it. Nests were established either in non-use areas or in areas of light to moderate grazing. These included both taller patch vegetation of draws and shorter uniform vegetation of open fields; greater hatching success occurred in the uniform areas. Important nest site species included crested wheatgrass (Agropyron cristatum), buckbrush (Symphoricarpos occidentalis), and sweet clover (Melilotus officinalis). Non-use grasslands, edges of heavily grazed pastures, and brushy draws were utilized extensively by broods.

## INTRODUCTION

This study was conducted to investigate and define the habitat requirements of the sharp-tailed grouse (*Pedioecetes phasianellus jamesi* Lincoln) during the reproductive season. A determination of specific requirements might indicate possible improvements for the land management systems currently in use in southwestern North Dakota, where approximately 1.1 million acres are administered by the U. S. Forest Service and Bureau of Land Management under the multiple-use concept. Therefore, new concepts, indicated by this and other studies, might be applied to this vast area resulting in a management system suitable for both livestock and grouse.

The study was conducted under a cooperative agreement between the North Dakota State Game and Fish Department, the U.S. Forest Service, and the University of North Dakota. The main objectives were (1) to determine the dispersal of sharptail hens from dancing grounds to nesting sites, (2) to determine the quality and quantity of selected nesting cover, (3) to determine the influence of intensive grazing management (rotation grazing) on grouse habitat and populations, and (4) to determine the home range, daily movement patterns, and cover preferences of sharptail broods.

## DESCRIPTION OF THE STUDY AREA

### Location

The study area is in northern Billings County, North Dakota west of Fairfield and northeast of Gorham (Figure 1). It is located in T.143N., R.99W., Sec. 16-21 and 28-33; T.143N., R. 100W., Sec. 13, 24, 25, and 36, and is bordered on the west by the Badlands and on the east by U.S. Highway 85. More than half of the 10,240 acres of the area is under U. S. Forest Service control.

### Native vegetation

The native grasslands of southwestern North Dakota commonly include dominant species from both the mixed and true prairie associations. Uplands are commonly occupied by needle-and-thread (*Stipa comata*), green needle (*S. viridula*), western wheatgrass (*Agropyron smithii*), sedges (*Carex spp.*), little bluestem (*Andropogon scoparius*), junegrass (*Koeleria cristata*), blue grama (*Bouteloua gracilis*), side-oats grama (*B. curtipendula*), fringed sage (*Artemisia frigida*), (Hanson and Whitman, 1938); the principal introduced species is crested wheatgrass (*A. cristatum*).

Trees and brush are mainly confined to draws or are found in close proximity to them. Woody species of importance include buckbrush, (*Symphoricarpos occidentalis*), buffalo berry (*Shepherdia argentea*), Juneberry (*Amelanchier alnifolia*), chokecherry (*Prunus virginiana*), western wild rose (*Rosa woodsii*), green ash (*Fraxinus pennsylvanica* var. *lanceolata*), cottonwood (*Populus deltoides*), poison ivy (*Rhus radicans*), currant (*Ribes spp.*), and Hawthorn (*Crataegus spp.*).

### Land-use

Sixty-six and three tenths percent of the area is grassland (Table 1); 33.7 percent is either small grain [cultivations] (15.3 percent), introduced pasture grasses (8.2 percent), or forage crops (10.2 percent). Wheat, barley, and oats are the important small grains; the latter two used primarily as feed grains. Beef cattle are raised under the cow-calf grazing system. Hay and alfalfa are utilized more extensively than native grasses for wintering cattle.





Figure 1. Location map of Gorham Study Area, North Dakota 1969-1970.

Table Land-use on 23,680 Acres on and Surrounding G... Study Area, Summer, 1968 (Bernhoft, 1969).

| Cover type          | Cultivated   |                  | Cover type     | Non-cultivated |                  |
|---------------------|--------------|------------------|----------------|----------------|------------------|
|                     | No. of acres | Percent of total |                | No. of acres   | Percent of total |
| Wheat               | 1,940.8      | 8.2              | Native pasture | 13,841.7       | 58.5             |
| Oats                | 288.1        | 1.2              | Native hay     | 882.6          | 3.7              |
| Barley              | 79.2         | 0.3              | Idle areas     | 190.4          | 0.8              |
| Rye                 | 26.4         | 0.1              | Trees          | 52.6           | 0.2              |
| Flax                | 5.4          | tr               | Road ditches   | 286.0          | 1.2              |
| Millet              | 9.1          | tr               | Fence rows     | 59.2           | 0.3              |
| Corn                | 73.9         | 0.3              | Farm steads    | 95.1           | 0.4              |
| Alfalfa             | 594.1        | 2.5              | Roads          | 105.4          | 0.4              |
| Mixed hay           | 1,756.5      | 7.4              | Soil bank      | 110.5          | 0.5              |
| Crested wheat grass | 1,683.8      | 7.1              | Field borders  | 9.0            | tr.              |
| Mixed pasture       | 249.2        | 1.1              | Marsh          | 16.3           | 0.1              |
| Summer fallow       | 1,284.6      | 5.4              | Open water     | 40.1           | 0.2              |
| TOTAL               | 7,991.1      | 33.7             |                | 15,688.9       | 66.3             |



# LITERATURE REVIEW

## Grouse habitat and range management

In South Dakota good prairie grouse habitat consisted of native pasture less than 50 percent utilized, unmowed hayland, trees, shrubs, and weeds; poor habitat included cultivated land, heavily utilized native grass, and mowed hayland. Areas with less than 50 percent good habitat had retrogressing densities, those with 50-60 percent had increasing densities, however, those with more than 70 percent did not have correspondingly higher densities (Janson, 1953; Podoll, 1955 and 1957; Frary, 1958; West, 1959). Janson (1953) recommended that sharptail habitat in South Dakota should consist of 74 percent grassland, one-third overgrazed or mowed, 14 percent small grains, 3 percent row crops, and 1.5 percent trees or brush.

In the Lake States, sharptails require more woody cover, nevertheless, they are still dependent upon open areas. Ideal prairie grouse habitat in this area should include 6 percent open grassland for dancing, feeding, and loafing; 50 percent mixed woodland and grassland for roosting, nesting, and feeding; and 44 percent woodland with small grassy openings for winter feeding and roosting, for resting, brood-rearing, and escape cover (Ammann, 1957). Grange (1948) defined the minimum unit of optimal sharptail range in Wisconsin as 2,000 acres. Uhlig and Hamor (1966) suggested that the minimum practical size of a sharptail habitat under ideal conditions would range from one square mile in forested to four square miles in grassland areas; five times as much area was required in poorer habitats.

The importance of brushy draws, shrubs, and woody vegetation has been well documented (Johnson, 1934; Marshall and Jensen, 1937; Hamerstrom and Hamerstrom, 1951b; Yocom, 1952; Janson, 1956), however, overbrowsing curtails woody vegetation (McKenzie, 1963). On the other hand, it has been demonstrated that the replacement of grassland in the Lake States by woody cover is more detrimental to grouse than overgrazing (Leopold, 1933; Hamerstrom and Hamerstrom, 1951a). Therefore, management entails maintenance of open areas by controlled burning, herbicides, and mechanical disturbances (Doll, 1955; Newman, 1959; Ammann, 1963; Mattson 1963).

Population densities of sharp-tailed grouse are directly related to the quality of the vegetation of grassland habitats. However, maintenance of normal plant cover or climax grassland is impossible under excessive grazing and trampling (Weaver, 1954). Correlations between the amount of grazing and erosion, and floral and faunal changes have been described by Smith (1940). Henderson (1964) found that heavily grazed native grasslands where annual utilization was 50-100 percent were unsuitable as grouse habitat; the greatest grouse populations were found where grazing was light or moderate, less than 50 percent. The adverse effect of increased grazing pressures and agricultural practices on sharptail populations have been well documented (Schrader and Erickson, 1944 in Minnesota; Buss and Dziedzic, 1955 in Washington; Janson, 1956 and Henderson, 1964 in South Dakota; Evans, 1961 in Alberta; Hamerstrom and Hamerstrom, 1961 in the Dakotas and Nebraska; Brown, 1963 and 1966a in Montana).

Uhlig and Hamor (1966) stated:

Proper range management that assures peak forage production and the maintenance of the best forage producing native grass species will provide adequate nesting, rearing, and roosting cover.

Mathison (1963) postulated that good range management is also good grouse management. Therefore, a very important criterion for sharp-tailed grouse management is the regulation of grazing to insure proper range management (Hart, Lee, and Low, 1950; Hamerstrom and Hamerstrom, 1961; Henderson, 1964).

#### Telemetry

Radio telemetry has been employed in the study of numerous wildlife species but was first applied to an upland game bird species, the ruffed grouse (Bonasa umbellus), by Marshall and Kupa (1963). Other investigators have utilized radio telemetry with various species of prairie grouse (Brown, 1965; Slade, Cebula, and Robel, 1965; Cebula, 1966; McEwen and Brown, 1966; Robel, 1966, Viers, 1967; Bernhoft, 1969; Robel et al., 1970). Telemetry also has been used in ring-necked pheasant studies by Kuck (1966). Bird activity has been successfully interpreted by changes in transmitter signals, i.e., the subject is stationary, walking, flying, feeding, or dusting (Marshall, 1963; Brown, 1966b; Kuck, 1966).

Although sharptails and prairie chickens carrying transmitters were thought to be more susceptible to predation, McEwen and Brown (1966) and Robel (1970) found them no more vulnerable than non-radio equipped birds. Studies of sharp-tails (Brown, 1965; Bernhoft, 1969), pheasants (Kuck, 1966), and prairie chickens (Cebula, 1966) also indicated that transmitters did not adversely affect normal activity. Kuck (1966) observed a copulation between a cock and a radio equipped hen pheasant.

#### Trapping techniques

Several trapping methods have been employed for capturing sharptail grouse during their reproductive season. The cannon-net described by Dill and Thornsberry (1950) was used to trap sharptails on dancing grounds (U.S. Fish and Wildlife Service, 1955; Peterle, 1956; Klett, 1957; Schwilling, 1960b; Kobriger, 1964; Brown, 1965; Bernhoft, 1969). The cannon-net has been mounted on the front of a vehicle to increase its mobility and effectiveness (Lacher and Lacher, 1964; Higby, 1969). Brown (1966b) and Bernhoft (1969) trapped sharptails on dancing ground with portable funnel traps.

Mist nets have been used to capture sharptails on dancing grounds (Evans, 1961), prairie chickens on booming grounds (Silvy and Robel, 1968), instrumented brooding female blue grouse (Dendragapus obscurus) (Schladweiler and Missehl, 1969), and female sharptails and broods (Artmann, 1969). Incubating sharptails have been taken with spring operated bow-net traps (Bernhoft, 1969) and with long handled hoop nets (Schwilling, 1960b; Brown, 1966b; Bernhoft, 1969). Hoop nets have been used to capture brooding sharptails (Brown, 1965) and incubating pheasants (Kuck, 1966). Brown (1966b) and Cebula (1966) netted sharptails by pulling a large net over them at night. Evans (1961) trapped incubating and brooding hens with funnel-type traps. Robel et al. (1970) captured greater prairie chickens on booming grounds with mist, bow, and cannon nets and incubating hens with a long handled hoop net.

#### Hen dispersal

According to Hamerstrom (1939) females travel only short distances from breeding grounds to nesting sites. Uhlig and Hamor (1966) stated that sharptails

usually nest within 1.0 mi. of dancing grounds and most authorities agree. Sharptail nests have been found within 0.5 mi. (Leopold, 1933), between 0.50 and 0.75 mi. (Symington and Harper, 1957), within 0.75 mi. (Klett, 1957), and within 1.0 mi. (Kobriger, 1964) of the nearest dancing ground.

#### Nesting characteristics

Sharptail hens must utilize available cover for nesting, regardless of its quality, because they ordinarily nest in the immediate vicinity of dancing grounds (Blus and Walker, 1966). In the Lake States most nests have been enclosed by heavy grass or concealed near or under logs, brush or at the base of small trees or bushes (Gross, 1930; Saunders, 1936; Baumgartner, 1939; Hamerstrom, 1939; Schrader and Erickson, 1944; Scott, 1947; Ammann, 1957; Bent, 1963; Evans, 1968).

In the Plains States Viehmeyer (1941) and Henderson (1964) considered that the most important basic needs of sharp-tailed grouse were nesting and roosting cover. A good nest site, according to Symington and Harper (1957), required an abundance of protective cover in the form of grasses and low shrubs and a readily available food supply. Schwilling (1960c) and Kobriger (1964) observed that prairie grouse nests were either in non-use or lightly grazed areas and that residual vegetation from the previous year's growth provided part of the required cover. Of 25 sharptail nests found by Bach (1943) none were on land extensively utilized by man, i.e., cultivated, closely mowed wild hay, or over grazed pasture. All were located on wild land partly used for pasture. Blus and Walker (1966) found that nests were usually in native prairie where little or no grazing had occurred; 73.3 percent were on range of excellent conditions, 23.4 percent on range of good condition, and 3.3 percent on range of fair condition; none were on range which was undergoing severe or extreme grazing. Fifteen and five tenths percent of the nests were located on the unused range which comprised less than 1 percent of the total area. Although Hart *et al.* (1952) and Blus and Walker (1966) found a few nests in stubble and alfalfa (*Medicago sativa*), the majority of nests begun in late April and early May were in patches of residual tall and medium height grasses, which persisted through the winter and spring (Janson, 1956), i.e., needlegrass (*Stipa* spp.) bluestem (*Andropogon* spp.), and wheatgrass (*Agropyron* spp.). Janson (1956) and Blus and Walker (1966) noted that nests were often located on north eastern exposures probably because of denser cover and a cooler and more moist microclimate. Sharp (1958) found that the body temperature of a ground nesting bird created a microclimate at the nest site which stimulated grass and forb growth at the periphery of the nest and hastened nest concealment.

The nesting season continues from late April to early July. More nests were begun during the peak of the dancing season in late April or early May (Hamerstrom, 1939; Uhlig and Hamor 1966; Evans, 1968). Captive sharptails laid their first egg on May 1 or 2 and reached a peak during the period May 22 to June 11; after the first week of July very few eggs were laid (McEwen, Knapp, and Hilliard, 1969). However, observers generally agree that the peak of hatching in the wild occurs during the first three weeks of June (Baumgartner, 1939; Bach and Stuart, 1942; Blus and Walker, 1966; Brown, 1967). The average number of eggs per sharptail clutch was approximately 12 (Baumgartner, 1939; Hamerstrom, 1939; Bach and Stuart, 1942; Schrader and Erickson, 1944; Scott, 1947; Blus and Walker, 1966; Brown, 1967). High fecundity has been found and fertility ranges up to 97 percent (Brown, 1967). Baumgartner (1939) did not note a change in the clutch size from year to year and Brown (1966b) did not find a correlation between clutch size and the age of hens.

Hamerstrom (1939), Cartwright (1944), Ammann (1957), and Blus and Walker (1966) have agreed that reproductive success depends on initial nesting efforts because prairie grouse are not persistent renesters. However, according to Brown (1966b) instrumented sharp-tails renested persistently; one hen renested twice in one summer. Successful renesting was measured by the spread of brood ages observed in August. Symington and Harper (1957) noted that the early destruction of first nests resulted in one or two additional attempts at renesting. Of 73 nests found by Blus and Walker (1966), 8.2 percent were renests; these accounted for only 2.3 percent of the chicks produced in 34 nests. Although a small sample was examined, it was observed that renests contain less eggs than initial nests. However, Brown (1966b) found no significant difference in the numbers of eggs in initial and renests. Sharptail nesting successes in North Dakota have been reported by Bach and Stuart (1942) and Bach (1943) as 68.0 percent of 25 and 86.2 percent of 29 nests, respectively. Ammann (1957) observed a hatching success of 44.0 percent of 330 Michigan prairie grouse nests. In a recent study of prairie grouse in the Nebraska Sandhills, Blus and Walker (1966) reported 49.3 percent hatch from 67 nests. They also noted a correlation between hatching success and range condition. Hatching success on an over-grazed and intensively mowed area was 40.0 percent of 35 nests and on an area which was less intensively utilized 61.3 percent of 31 nests hatched; one nest was unaccounted for. Uhlig and Hamor (1966) stated that in average years approximately 50 percent of the sharptail nests are successful.

The reproductive success may be related to the weather. Janson (1955) and Hamerstrom and Hamerstrom (1961) agreed that the greatest percentage of success occurred during relatively cool, wet summers; warm, dry weather was unfavorable to reproduction. During the drought periods of 1910-11, 1925-27, and 1934-37 there were disastrous reductions in sharptail nesting success (Janson, 1955). Zwickel (1958) suggested that weather affects blue grouse reproduction, i.e., years with warmer, wetter Mays enhanced the cover; dryer, warmer Junes favored broods. Young prairie grouse (Cartwright, 1944; Schwartz, 1945; Scott, 1947) and pheasant chicks (MacMullan and Eberhardt, 1953) are most vulnerable to weather extremes, particularly wet cold periods. A 50 percent mortality occurs during the first 3 weeks of life in early sharptail broods (Brown, 1967). Four or at the most five, chicks from a clutch of 12 eggs can be expected to reach maturity (Symington and Harper, 1957).

#### Predation

The mechanism of staggering nesting attempts guarantees the survival of short lives species because it insures that the total hatch would not be vulnerable to a general disaster at any one period of the reproductive season (Cartwright, 1944; Symington and Harper, 1957). However, Darrow (1945) indicated that the relative abundance of buffer species will determine the predatory pressure on ruffed grouse and their nests. Thus, predation must have been an important factor in an annual sharptail mortality of 60 to 70 percent (Henderson, 1964). Brown (1966b) reported that coyote predation on nests and nesting hens was a major factor limiting total sharptail reproduction.

#### Brood Habitat

The influence of season, intensity, and type of grazing on the structure and composition of vegetation during the brood season have been discussed by Craddock and Forsling (1938) and Miegler (1962). In early summer, sharptail broods utilized a variety of cover, generally within the immediate vicinity of the nest (Ammann, 1957; Klett, 1957; Brown, 1966b). Edminster (1954), Peterle (1954), and Ammann (1957) reported that broods moved to denser cover some dis-



tance from the nest site later in the summer, thus increasing brood range. However, Uhlig and Hamor (1966), Evans (1968), and Bernhoft (1969) have stated that summer brood territory rarely exceeds an area one-half mi. in diameter. Hamerstrom (1963) investigated 207 locations of 193 sharptail broods in the pine barrens of Wisconsin and found 80 percent in open cover types, 14 percent in edge habitats, and 5 percent more than 50 yards into woodland. Grassland with shrubs was determined to be the most important cover. Baumgartner (1939) found 17 broods in open uplands within 0.67 mi. of the nearest dancing ground with but one exception. Jackson (1963) studied 44 sharptail broods and found 67 percent of them in ungrazed and 33 percent in grazed areas; 29 were in native grasses, 16 in exotic grasses and 10 in annual weeds. Natural woody cover along draws offered shelter and shade to broods in the summer (Henderson, 1964).

Kobriger (1964) observed that broods roosted at night in upland sandhills on the Valentine Refuge in Nebraska and spent days in the mowed wetland meadows. During the day any available shade, including trees, windrows, unmowed areas, and other patches of vegetation, were utilized. He also reported that broods abandoned wetland meadows following mowing. Brown (1966b) noted the concentration of broods following the early dessication of grasses and forbs in a year of low rainfall; occasional grouping of broods or broods and non-brooding hens were also observed. The daily sequence of hen-brood activity consisted of night-roosting, morning feeding, day resting, and afternoon feeding; high temperatures, wind, and rain disrupted this normal cycle.

#### Vegetation analysis

There is a paucity of information concerning methods and techniques employed in evaluating nesting and brooding cover. Emlen (1956) developed a method to describe and compare avian habitats based on their measurable features, including characteristics of the vegetation, substrata, and surrounding area. Vegetational height was considered the most important factor by Lack (1933) who employed this to differentiate various habitats. Gross cover requirements of ruffed grouse were described by Polderboer (1942) and Dorney (1959). Evans and Gilbert (1969) systematically assigned a numerical value to rate individual habitat components, including winter food, protective cover, nesting cover, area (grassland), shrub cover, topography, soil, and water. In aggregate these ratings indicated the potential value of a specific range. Separately, they indicated those range components which required improvement. Jones (1960) recorded the physiognomic life-form types at prairie chicken flushing sites under three principal divisions: woody plants, half-shrubs, and herbs, with subdivisions according to height of vegetation.

Various types of equipment have been used in cover analysis, such as, photoelectric equipment to measure density (Allard, 1947; Sather, 1950) and photography for cover evaluation (Bump, 1950; Mosby, 1963). Wight (1939), Kobriger (1964), Jones (1968), and Bernhoft (1969) employed density boards to visually measure the cover of deer habitat, grouse breeding ground cover, grouse coverts, and grouse nesting cover, respectively. Jenkins (1961) and Watson (1964) described techniques to evaluate specific cover types for Hungarian partridge (Perdix perdix perdix) and red grouse (Lagopus lagopus scoticus), respectively. Both use visibility of models of the species being studied as the criterion for rating the particular cover.

Bue, Blankenship, and Marshall (1955) evaluated waterfowl nesting cover using height-density class rating. As grazing increased, there was a concomitant reduction in rating and total number of nests. Bernhoft (1969) adapted the clipped hoop method of range analysis to evaluate nesting cover for sharptails. The

degree of use guide described by Dyksterhuis (1944) was used by Schwilling (1960a) to determine preferred nesting cover of prairie grouse. This was based on livestock utilization in the vicinity of nest sites.

#### METHODS AND MATERIALS

##### Trapping and marking

Field work continued from April 15 through August 20, 1969 and from April 1 through August 26, 1970. Work began with inventory of dancing grounds to determine areas of female concentrations and activity in order to place traps in the most advantageous locations for capturing birds. The areas were prebaited with corn and wheat before single three-funnel (6' X 6' X 3') walk-in traps were set up on dancing grounds 1, 4, and 6 in 1969. Ground 4 was also trapped with a double, Miller-improved cannon-net assembly (75' X 50') in 1969. In 1970 a double, Dill recoilless cannon-net assembly (60' X 40') was used on grounds 2, 4, and 5. All trapped birds were weighed, sexed, and banded with numbered, aluminum, open-end type leg bands and color-coded celluloid leg bands to designate the dancing ground from which the birds were trapped.

A long handled hoop net and a spring operated bow-net trap were used to recapture incubating hens for battery replacement and weighing. One incubating hen was taken from her nest by hand and one brooding hen was recaptured with a bumper mounted mobile cannon-net (50' X 40'). The long handled hoop net was also utilized to capture young brood members during the summer.

##### Telemetry

Sharptail hens were equipped with radio transmitters according to the technique of Brander (1968) and released. The battery leads were connected, soldered and the entire battery component was taped and then waterproofed with dental acrylic, liquid-tape (G. C. Electronics, Division of Hydrometals, Inc., Rockford, Illinois). Hens were located with a portable receiver and hand-directional antenna. Telemetry equipment (Sidney L. Markusen, Electronics Specialties, Esko, Minnesota) was similar to that of Bernhoff (1969) with two exceptions. A second portable receiver was employed as a back-up unit and only the 28 inch, portable, hand-directional antenna was used; the stationary yagi antenna was abandoned.

##### Dispersal of hens

Radio equipped hens were released near the dancing ground trap sites. They were monitored for dispersal and nest selection. Dispersal distances were measured on a straight line from the dancing ground to the nest sites.

##### Vegetation at nest sites

Vegetation at 22 nesting sites was sampled using the clipped hoop method of range analysis (U.S. Department of Agriculture, Forest Service, 1969). Three nest sites were not clipped as the cover had changed drastically by being mowed, plowed, or destroyed by hail. Herbage within two hoops, each subtending a 0.96 square foot area, was clipped at opposite edges of the nest. This was identified, sacked, and air dried for subsequent weight determination. Clippings of two 1969

nests were made in early August following seasonal growth of the vegetation. Nine of twelve 1969 nests were clipped during the spring of 1970 on or near the dates the hens occupied the sites the previous year. All 1970 nests were clipped as early as possible in the reproductive season after hatching, predation, or desertion. Production was measured in pounds per acre and percentage of each plant species. The former was computed by multiplying the weight in grams of a species found within the two hoops by 50.

As soon as nests were located, a 2 X 1 foot cover board, horizontally marked with alternating one inch white and black strips was temporarily placed at two adjacent edges of each nest to visually analyze the cover. This board was removed after the observations were made. Height and density of vegetation was determined by the following two procedures: (1) observations of the number of strips completely covered by vegetation were made giving a 100 percent density-height factor; and (2) estimations of the maximum height of the vegetation were made from the percent of the board obscured by the taller vegetation above the 100 percent density-height factor (Bernhoft 1969).

Paced line transects were run in the vicinity of seven nest sites in pastured areas. Vegetative and soil conditions were rated according to a five division scale of excellent, good, fair, poor and very poor; vegetative and soil trends were also noted (U.S. Department of Agriculture, Forest Service 1969).

#### Brood movement and cover selection

For each brood seen during the summer, the location, adjacent land-use, vegetative cover height and density, and species composition were recorded. In addition to these occasional sightings, intensive observations were made of broods of radio equipped hens. Daily movements and activities of these broods were determined with up to three individual contacts per day. Occasional days with dawn-to-dusk checking at 15 minute intervals supplemented the data collected by the normal daily checks. Instrumented hens and broods were located visually without flushing to determine vegetation preferences. Brood survival was determined by periodic counts on those days when broods were accidentally flushed.

## RESULTS

#### Trapping and marking

Sharptails were trapped on the five active dancing grounds within the study area (Table 2). Of 62 sharptails captured during the study, 36 were males and 26 were females. Weight data for these birds are summarized in Table 3. In 1969, ten males were cannon-netted on ground 4, one was caught on ground 1 with a walk-in trap, and four were caught in a walk-in on ground 6. The male from ground 1 and one from ground 6 had been banded on their respective grounds the previous spring. Two young of the year males were hoop netted during the summer. In 1970, 19 males were cannon-netted on three grounds, four on ground 2, two on ground 4, and thirteen on ground 5. One male from ground 4 had been banded the previous spring on the same ground and one male from ground 5 had been banded in the spring of 1967 on ground 5, then located one-half mile to the east. Of 17 females taken in 1969, 16 were cannon-netted on ground 4 and one young of the year female was hoop netted during the summer. In 1970 eight females were cannon-netted on three grounds, five on ground 2, one on ground 4, and two on ground 5. During the summer one young of the year female was hoop netted. A summary of all trapping is presented in Table 4.

Table 2. Male Sharp-tailed Grouse Census, Gorham Study Area, North Dakota, Spring 1966-1970.

| Dancing Ground | Number of Males   |                   |           |           |           |
|----------------|-------------------|-------------------|-----------|-----------|-----------|
|                | 1966 <sup>a</sup> | 1967 <sup>b</sup> | 1968      | 1969      | 1970      |
| 1              | 8                 | 16                | 16        | 4         | 2         |
| 2              | 17                | 19                | 14        | 11        | 9         |
| 3              | 9                 | 4                 | 0         | 0         | 0         |
| 4              | 23                | 23                | 24        | 23        | 16        |
| 5              | 29                | 31                | 11        | 9         | 15        |
| 6              | --                | --                | 24        | 10        | 10        |
| Total          | <u>86</u>         | <u>93</u>         | <u>89</u> | <u>57</u> | <u>52</u> |

<sup>a</sup>North Dakota State Game and Fish Department

<sup>b</sup>1967 and 1968 Data (Bernhoft, 1969)

Table 3. Weights of Adult and Immature Sharp-tailed Grouse, Gorham Study Area, North Dakota, 1969-1970.

| Class            | Number | Weight (grams) | Range     | Mean |
|------------------|--------|----------------|-----------|------|
| Adult males      | 34     | 31,033         | 794-1,077 | 913  |
| Adult females    | 24     | 20,097         | 709-1,134 | 837  |
| Immature males   | 2      | 369            | 156-213   | 184  |
| Immature females | 2      | 737            | 255-482   | 368  |



Table 4. Sharp-tailed Grouse Trapping Methods and Success, Gorham Study Area, North Dakota, 1969-1970.

| Ground No.           | Method            | Trap Days |      | Males          |                 | Females |      | Total |
|----------------------|-------------------|-----------|------|----------------|-----------------|---------|------|-------|
|                      |                   | 1969      | 1970 | 1969           | 1970            | 1969    | 1970 |       |
| 1                    | Funnel Walk-in    | 14        | --   | 1 <sup>c</sup> | --              | 0       | --   | 1     |
| 2                    | Cannon-net        | --        | 6    | --             | 4               | --      | 5    | 9     |
| 4                    | Funnel Walk-in    | 14        | --   | 0              | --              | 0       | --   | 0     |
|                      | Cannon-net        | 2         | 1    | 10             | 2 <sup>d</sup>  | 16      | 1    | 29    |
| 5                    | Cannon-net        | --        | 6    | --             | 13 <sup>e</sup> | --      | 2    | 15    |
| 6                    | Funnel Walk-in    | 14        | --   | 4              | --              | 0       | --   | 4     |
| Nest <sup>a</sup>    | Hoop net          | --        | --   | --             | --              | 5       | 2    | 7     |
|                      | Nest trap         | --        | --   | --             | --              | 1       | 4    | 5     |
|                      | Mobile Cannon-net | --        | --   | --             | --              | --      | 1    | 1     |
| General <sup>b</sup> | Hoop net          | --        | --   | 2              | --              | 1       | 1    | 4     |
| Total                |                   | 44        | 13   | 17             | 19              | 23      | 16   | 75    |

<sup>a</sup>All radioed hens recaptured for battery change

<sup>b</sup>Immature birds

<sup>c</sup>1 banded in 1968

<sup>d</sup>1 banded in 1969

<sup>e</sup>1 banded in 1967

Of six returns (4 males and 2 females) in the fall of 1969, five, including both females, were banded during the previous spring and one was banded during the spring of 1968. One return in 1970 had been banded during the previous spring. All birds were killed within one mile of their home grounds with the average distance being approximately one-half mile.

#### Telemetry

Of 18 females trapped, radio-equipped, and released, eleven were taken on ground 4, five on ground 2, and two on ground 5. Table 5 presents the weight changes of six radio-equipped hens between the spring capture and the mid-incubation recapture dates. One young female brood member was captured, radio-equipped, and released during the summer but contact was lost the next day.

The effective range of transmitters was 0.75 mi., however, under ideal conditions, the range was up to 2.0 mi. Topography, density of vegetation, and location and activity of birds affected normal transmission and reception.

Battery life varied throughout the study. Of 31 transmitters operational for 12 to 110 days, averaging 53 days, only four were lost unexpectedly. These were probably due to battery failures.

#### Dispersal of hens

Eighteen radio-equipped females were monitored from dancing grounds to nesting areas to provide dispersal information (Figure 2 and Table 6). Table 6 also presents the results of 17 initial nests and eight re-nests. In 1969 ten hens were monitored for 925 days and in 1970 eight hens were monitored for 714 days. Radio monitoring data and the final disposition of each hen is summarized in Table 7.

Table 5. Comparison of Initial Spring and Mid-incubation Weights<sup>a</sup> of Six Sharp-tailed Grouse Hens, Gorham Study Area, North Dakota, 1969-70.

| No.                 | Spring  |        | Mid-incubation |        | Weight Loss <sup>c</sup> |
|---------------------|---------|--------|----------------|--------|--------------------------|
|                     | Date    | Weight | Date           | Weight |                          |
| 1-69                | 4-23-69 | 907    | 6-11-69        | 822    | 85                       |
| 2-69                | 4-23-69 | 822    | 6-4-69         | 765    | 57                       |
| 4-69                | 4-23-69 | 765    | 6-10-69        | 680    | 85                       |
| 7-69                | 4-23-69 | 879    | 6-11-69        | 822    | 57                       |
| 2-70                | 5-4-70  | 879    | 6-17-70        | 737    | 142                      |
| 10 <sup>i</sup> -70 | 4-30-70 | 794    | 6-17-70        | 709    | 85                       |

<sup>a</sup>In grams

<sup>b</sup>Mean weight loss 85

<sup>i</sup>Intermittent signal channel




Figure 2. Hen Dispersal from Dancing Grounds, Gorham Study Area, North Dakota, 1969-1970.

Table 6. Sharp-tailed Grouse Hen Dispersal and Nesting, Gorham Study Area, North Dakota, 1969-1970.

| Channel             | Dispersal (Miles) |                   | Direction<br>from<br>dancing<br>ground | Nesting<br>results              |
|---------------------|-------------------|-------------------|--|---------------------------------|
|                     | Maximum           | Nest <sup>a</sup> |  |                                 |
| 1-69                | 1.0               | 0.9               | W                                      | Egg predation                   |
| 2-69                | 0.8               | 0.8               | SSW                                    | Hatched                         |
| 3-69                | 2.0               | 2.0               | NNE                                    | Hen predation                   |
| 4 <sup>a</sup> -69  | 0.9               | 0.9               | WSW                                    | Egg predation                   |
| 4 <sup>b</sup> -69  | 1.5               | 1.2               | WSW                                    | Hen and egg predation           |
| 5-69                | 0.9               | 0.8               | SW                                     | Hatched                         |
| 6-69                | 1.1               | 1.1               | NW                                     | Hatched                         |
| 7-69                | 0.9               | 0.9               | W                                      | Egg predation                   |
| 8 <sup>a</sup> -69  | 0.8               | 0.4               | ENE                                    | Egg predation                   |
| 8 <sup>b</sup> -69  | 0.8               | 0.7               | NE                                     | Hatched                         |
| 10 <sup>a</sup> -69 | 1.1               | 0.8               | E                                      | Egg predation                   |
| 10 <sup>b</sup> -69 | 1.1               | 0.7               | ENE                                    | Hatched                         |
| 1 <sup>a</sup> -70  | 0.5               | 0.5               | WSW                                    | Nest predation                  |
| 1 <sup>b</sup> -70  | 0.6               | 0.6               | NNE                                    | Hatched                         |
| 2-70                | 0.5               | 0.3               | W                                      | Hatched                         |
| 3 <sup>a</sup> -70  | 0.9               | 0.2               | N                                      | Nest predation                  |
| 3 <sup>b</sup> -70  | 0.4               | 0.3               | ENE                                    | Hen lost (mammalian predation?) |
| 6-70                | 1.7               | 1.7               | SSW                                    | Hatched                         |
| 8-70                | 0.5               | 0.5               | NW                                     | Hatched                         |
| 10 <sup>a</sup> -70 | 0.4               | 0.3               | NNW                                    | Nest predation                  |
| 10 <sup>c</sup> -70 | 0.4               | 0.4               | NW                                     | Abandoned                       |
| 10 <sup>b</sup> -70 | 0.4               | 0.4               | NW                                     | Abandoned                       |
| 12 <sup>a</sup> -70 | 0.8               | 0.5               | NW                                     | Nest predation                  |
| 12 <sup>b</sup> -70 | 0.3               | 0.3               | NNW                                    | Hatched                         |
| 10 <sup>i</sup> -70 | 0.8               | 0.8               | S                                      | Hatched                         |

<sup>a</sup>Initial nest

<sup>b</sup>Renest

<sup>c</sup>Nest bowl

<sup>d</sup>Mean distance 0.8

<sup>i</sup>Intermittent signal channel

Table 7. Telemetric Study and Final Disposition of 18 Sharp-tailed Grouse Hens, Gorham Study Area, North Dakota, 1969-1970.

| Channel No.         | Date                 |                     | Monitoring Discontinued | Total Days Monitored | Battery Life |        | Disposition                        |
|---------------------|----------------------|---------------------|-------------------------|----------------------|--------------|--------|------------------------------------|
|                     | Instrumented Initial | Instrumented Second |                         |                      | Initial      | Second |                                    |
| 1-69                | 4-23                 | 6-11                | 8-20                    | 121                  | 50           | 71     | Study terminated                   |
| 2-69                | 4-23                 | 6-3                 | 9-20                    | 152                  | 42           | 110    | Hen collected                      |
| 3-69                | 4-23                 | -                   | 7-18                    | 80                   | 80           | -      | Avian predation                    |
| 4-69                | 4-23                 | 6-10                | 6-22                    | 61                   | 49           | 12     | Mammalian predation                |
| 5-69                | 4-23                 | -                   | 6-12                    | 44                   | 44           | -      | Signal lost                        |
| 6-69                | 4-23                 | 6-2                 | 7-23                    | 93                   | 41           | 52     | Signal lost                        |
| 7-69                | 4-23                 | 6-11                | 8-20                    | 121                  | 50           | 71     | Study terminated                   |
| 8-69                | 5-1                  | -                   | 8-8                     | 100                  | 100          | -      | Signal lost                        |
| 9-69                | 4-23                 | -                   | 4-30                    | 7                    | 7            | -      | Signal lost                        |
| 10-69               | 5-1                  | 7-6                 | 9-20                    | 146                  | 79           | 67     | Hen collected                      |
| Subtotal            |                      |                     |                         | 925                  |              |        |                                    |
| 1-70                | 5-16                 | 7-9                 | 8-11                    | 87                   | 54           | 33     | Avian predation                    |
| 2-70                | 5-5                  | 6-17                | 8-25                    | 112                  | 43           | 69     | Study terminated                   |
| 3-70                | 5-5                  | -                   | 7-8                     | 64                   | 64           | -      | Signal lost (Mammalian predation?) |
| 6-70                | 5-5                  | 6-16                | 7-8                     | 64                   | 42           | 22     | Avian predation                    |
| 8-70                | 5-12                 | 6-24                | 8-8                     | 88                   | 43           | 45     | Mammalian predation                |
| 10-70               | 5-5                  | 7-9                 | 8-13                    | 100                  | 65           | 35     | Mammalian predation                |
| 12-70               | 5-5                  | 7-21                | 8-25                    | 112                  | 77           | 35     | Study terminated                   |
| 10 <sup>i</sup> -70 | 4-30                 | 6-17                | 7-25                    | 87                   | 49           | 38     | Mammalian predation                |
| Subtotal            |                      |                     |                         | 714                  |              |        |                                    |
| Grand total         |                      |                     |                         | 1639                 |              |        |                                    |

<sup>i</sup>Intermittent signal channel

Channel 1-69, instrumented April 23, was monitored for 121 days; the hen's recorded activity centered around a shallow draw containing buckbrush 0.9 mi. west of dancing ground 4. A nest, containing eleven eggs, was discovered in this area on June 9. This nest was destroyed June 20 and the hen moved to a draw 1.5 mi. southwest of the dancing ground and remained in this vicinity for the rest of the summer.

Channel 2-69, instrumented April 23, was tracked for 156 days. This hen remained close to a draw bordering a crested wheatgrass field in which her nest of thirteen eggs was discovered May 13, 0.8 mi. southwest of dancing ground 4. Eight of the final total of fifteen eggs hatched June 13.

On April 30, channel 3-69, was equipped with a radio and was followed for 56 days. Until May 19 the hen remained in a draw 1.4 mi. northeast of dancing ground 4. An additional dancing ground later discovered at this location could have been attended by this female. On May 20, the hen moved 0.6 mi. further northeast to a deep draw in which the nest was discovered May 30. It was located in a buckbrush patch and contained seven eggs. On June 14, the hen was missing from the nest and the radio signal could not be heard. Several feathers were found at the nest site indicating possible predation on the hen although the nest was intact. On July 18 the remains of the hen were found on a high butte 0.7 mi. west of the nest site. It was assumed that an avian predator killed this particular sharp-tail.

Channel 4-69 was released April 23 and followed for 61 days. This hen's nest was located on a crested wheatgrass hilltop 0.9 mi. west southwest of dancing ground 4. Nest predation occurred May 20, and the hen moved west across a road to a sweet clover field. The hen re-nested in this field, 1.2 mi. west southwest of the dancing ground. The re-nest, found June 9, and the hen was preyed upon June 22; the bird remains were found approximately 150 yards west of the nest.

Channel 5-69, initiated April 30, was monitored for 44 days. This hen occupied the same draw and crested wheatgrass field as channel 2. Channel 5's nest was discovered in the crested wheatgrass field 0.8 mi. southwest of dancing ground 4 on May 18 and contained nine eggs. The completed clutch contained 11 eggs of which only eight hatched on June 11. The entire brood was killed apparently by cold (chill factor 14F.) during a prolonged and unsuccessful early morning attempt to capture the hen to replace an inoperative transmitter.

Channel 6-69 was followed for 93 days after release on April 23. The hen located in a draw 1.1 mi. northwest of dancing ground 4 and her nest of 13 eggs was found under a buffalo berry bush on the slope of the draw May 12. The entire clutch hatched on June 6, coincidental with a thunderstorm with large hail which resulted in the loss of the entire brood. The hen remained in the area for the entire summer.

Instrumented April 23, channel 7-69 was tracked for 121 days. The hen located along a creek 0.9 mi. west of dancing ground 4 and nested on the side of a draw in buckbrush 50 yards from channel 1-69's nest. The nest containing eight eggs was found on June 5. Predation destroyed the clutch of ten eggs on June 23. The hen remained in the draw and pasture area 1.0 mi. northeast of the nest site and 0.75 mi. northwest of the dancing ground. This area was also used by channel 6-69 and the two birds were occasionally located together.

Channel 8-69, released May 1, was followed for 100 days. She located in a pasture containing shallow draws and ranged up to 0.8 mi. east of dancing ground 4. Her nest containing three eggs was located May 12 in the bottom of a draw containing smooth brome (Bromus inermis), 0.4 mi. northeast of the dancing ground. Nest predation destroyed 11 eggs on June 6. The bird renested 0.7 mi. northeast of the dancing ground in the same shallow draw. This nest, which was located June 23, was under a single large sweet clover plant in a smooth brome-alfalfa field. A total of nine eggs hatched July 17.

Channel 9-69 was lost the day following release, April 24. She was recannotted on dancing ground 4 April 30, but was subsequently lost again with no data collected.

Channel 10-69 was released May 1 and followed for 146 days. The hen located in the same pasture and shallow draw area as channel 8-69 and ranged up to 1.1 mi. east of dancing ground 4. Her nest of 11 eggs was found May 12 in the edge of dense buckbrush in a shallow draw 0.8 mi. east of the dancing ground. The total clutch of 13 eggs was destroyed by a predator June 6. Renesting took place and a completed nest of nine eggs was located June 23. This was in sparse buckbrush on a flat adjacent to the draw where her first nest was located and was 0.7 mi. east northeast of the dancing ground. On July 20, six of eleven eggs hatched.

Channel 1-70 was instrumented May 16 and followed for 87 days. She centered her activities around an aspen (Populus tremuloides) draw 0.5 mi. west of dancing ground 5. Her nest of 11 eggs was found May 29 under a buffalo berry bush at the edge of this draw. Nest predation occurred June 4. A second nest containing nine eggs was discovered June 18 on a hillside between two side draws 100 yards from the first nest. Seven of the completed clutch of 10 eggs hatched July 15.

Channel 2-70, released May 5, was monitored for 112 days. Two days following release a copulation was observed between this hen and a male on dancing ground 2. She located in an alfalfa field west across a draw from the dancing ground. Thirteen eggs were found in her nest in the alfalfa May 27, 12 of the final clutch of 14 eggs hatched June 25.

Channel 3-70, released May 5, was tracked for 64 days. Her initial nest of 11 eggs was found May 29 in quackgrass (Agropyron repens) and annual weeds north across a draw from dancing ground 2. Nest predation occurred on June 2. A renesting attempt produced 14 eggs in a nest located 0.3 mi. east northeast of the dancing ground in quackgrass and buckbrush. On July 9 the hen was missing from the nest and never located again. Mammalian predation may have taken place in the heavy nesting cover because numerous feathers were found near the nest site.

Channel 6-70 was monitored for 64 days after release May 5. On May 29, a nest containing 15 eggs was found 1.7 mi. south southeast of dancing ground 2 in a non-use area on Cropland Adjustment Program land. The entire clutch hatched June 22.

Channel 8-70, instrumented May 12, was followed for 88 days. A nest of 13 eggs was found May 29, 0.5 mi. northwest of dancing ground 5 in the crested wheatgrass used previously by channel 2-69 and 5-69. All eggs hatched June 22.



Channel 10-70 was released May 5 and followed for 100 days. Her first nest of six eggs, located 0.3 mi. north northwest of dancing ground 2, was found June 8 and destroyed by a predator June 14. On June 22 she was flushed from a scratched out depression that appeared to be the beginning of a nest bowl 0.4 mi. northwest of the dancing ground. Abandonment of the nest bowl apparently occurred because five eggs were found in a new nest 20 yards from the depression on June 29. This nest of 8 eggs was abandoned on July 9 after the hen was reinstrumented prematurely during the first few days of incubation to replace a non-functional transmitter. The hen remained in the draw and pasture 0.2 to 0.7 mi. northwest of the dancing ground for the remainder of the summer.

Channel 12-70, released May 4, was tracked for 112 days. The initial nest of 9 eggs was found 0.5 mi. northwest of dancing ground 2 on May 24. A clutch of 16 eggs had been laid before nest predation occurred June 14. The product of renesting was a nest of 7 eggs located June 28 in a crested wheatgrass area 0.3 mi. north northwest of the dancing ground. The entire clutch hatched on July 23 during a period of numerous thundershowers which killed the entire brood. The hen remained in the vicinity of the dancing ground the entire summer.

Channel 10<sup>i</sup>-70 was released April 30 and monitored for 87 days. On May 31 a nest with a clutch of 15 eggs was found at the edge of a wooded draw 0.8 mi. south of dancing ground 4. Fourteen eggs hatched June 24.

#### Vegetation at nest sites

Species composition of vegetation at nine 1969 sites and thirteen 1970 nest sites is presented in Table 8-14, respectively. The cover in proximity to three 1969 nests not analyzed consisted of a dense sweet clover field, a smooth brome-alfalfa field with a single sweet clover plant over the nest site, and the edge of a dense buckbrush patch in a draw.

The results of the visual analyses of all the nesting covers using a 1' x 2' cover board are shown in Table 15. Nineteen nests were located in pastures with light to moderate grazing, three in haylands, and three in non-use areas.

When the initial nests were found residual cover comprised from 10 to 90 percent of all cover present. In re-nests residual cover varied from 0 to 60 percent of all cover.

The vegetative condition of the vicinity of four of seven nests, as determined by paced line transects, were excellent; the other three were rated as good. In six of seven cases the vegetation trend was up; in all cases the soil trend was up.



Table 8. Species Composition and Amount of Vegetation at Channel 4-69 and 10-69 Nest Sites, Gorham Study Area, North Dakota. Vegetation Sampled by Clipped Hoop Method, Fall 1969 and Spring 1970.

| Species                            | Channel<br>Date | 4 <sup>a</sup><br>8-10-69 |      | 4 <sup>a</sup><br>5-6-70 |      | 10 <sup>b</sup><br>8-10-69 |       | 10 <sup>b</sup><br>6-20-70 |       |
|------------------------------------|-----------------|---------------------------|------|--------------------------|------|----------------------------|-------|----------------------------|-------|
|                                    |                 | lbs./acre                 | %    | lbs./acre                | %    | lbs./acre                  | %     | lbs./acre                  | %     |
| <u>Agropyron cristatum</u>         |                 | 1372.0                    | 68.0 | 740.5                    | 39.3 |                            |       |                            |       |
| <u>Agropyron smithii</u>           |                 |                           |      |                          |      | 122.0                      | 6.1   | 170.0                      | 4.3   |
| <u>Bromus japonicus</u>            |                 |                           |      |                          |      |                            |       | 58.0                       | 1.5   |
| <u>Buchloe dactyloides</u>         |                 | 11.0                      | 0.5  |                          |      |                            |       |                            |       |
| <u>Dantonion spp.</u>              |                 |                           |      |                          |      | 8.0                        | 0.4   |                            |       |
| <u>Koeleria cristata</u>           |                 | 185.0                     | 9.2  | 206.0                    | 10.9 | 72.5                       | 3.6   |                            |       |
| <u>Muhlenbergia cuspidata</u>      |                 |                           |      |                          |      |                            |       | 28.0                       | 0.7   |
| <u>Panicum capillare</u>           |                 |                           |      | 112.0                    | 5.9  |                            |       |                            |       |
| <u>Poa pratensis</u>               |                 |                           |      |                          |      | 132.5                      | 6.6   |                            |       |
| <u>Stipa comata</u>                |                 | 400.5                     | 19.9 | 805.5                    | 42.7 |                            |       |                            |       |
| <u>Stipa viridula</u>              |                 |                           |      |                          |      | 125.5                      | 6.3   | 180.0                      | 4.5   |
| <u>Achillea lanulosa</u>           |                 |                           |      |                          |      | 29.5                       | 1.5   | 190.0                      | 4.8   |
| <u>Artemisia frigida</u>           |                 | 10.5                      | 0.5  |                          |      | 112.5                      | 5.6   |                            |       |
| <u>Carex spp.</u>                  |                 |                           |      |                          |      | 125.0                      | 6.3   | 120.0                      | 3.0   |
| <u>Crepis occidentalis</u>         |                 |                           |      |                          |      | 6.0                        | 0.3   |                            |       |
| <u>Lotus americanus</u>            |                 |                           |      |                          |      | 27.5                       | 1.4   |                            |       |
| <u>Mamillaria spp.</u>             |                 | 34.5                      | 1.7  |                          |      |                            |       |                            |       |
| <u>Melilotus officinalis</u>       |                 |                           |      |                          |      | 73.5                       | 3.7   |                            |       |
| <u>Rosa arkansana</u>              |                 |                           |      |                          |      | 19.5                       | 1.0   | 120.0                      | 3.0   |
| <u>Symphoricarpos occidentalis</u> |                 |                           |      |                          |      | 1117.0                     | 55.9  | 2673.0                     | 67.2  |
| <u>Taroxicum officinale</u>        |                 |                           |      |                          |      | 8.0                        | 0.4   |                            |       |
| <u>Tragapogon dubius</u>           |                 |                           |      |                          |      |                            |       | 174.0                      | 4.4   |
| Unclassified forbs                 |                 | 3.0                       | 0.1  | 20.5                     | 1.1  | 19.5                       | 1.0   | 263.0                      | 6.6   |
| Total                              |                 | 2016.5                    | 99.9 | 1884.5                   | 99.9 | 1998.5                     | 100.1 | 3976.0                     | 100.0 |

<sup>a</sup>Initial nest

<sup>b</sup>Renest

9. Species Composition and Amount of Vegetation at Channel 1-69, 3-69, and 7-69 Nest Sites, Gorn  
Study Area, North Dakota. Vegetation Sampled by Clipped Hoop Method, Spring 1970.

| Species                            | Channel<br>Date | 1         |       | 3         |       | 7         |       |
|------------------------------------|-----------------|-----------|-------|-----------|-------|-----------|-------|
|                                    |                 | 5-26-70   |       | 5-26-70   |       | 5-28-70   |       |
|                                    |                 | lbs./acre | %     | lbs./acre | %     | lbs./acre | %     |
| <u>Agropyron smithii</u>           |                 | 22.0      | 0.4   | 5.0       | 0.1   | 76.0      | 1.9   |
| <u>Koeleria cristata</u>           |                 |           |       | 65.0      | 1.3   |           |       |
| <u>Poa pratensis</u>               |                 | 30.0      | 0.5   |           |       | 7.5       | 0.2   |
| <u>Stipa comata</u>                |                 |           |       | 13.5      | 0.3   |           |       |
| <u>Stipa viridula</u>              |                 | 2680.0    | 46.9  | 1368.0    | 27.2  | 514.0     | 12.9  |
| <u>Achillea lanulosa</u>           |                 |           |       |           |       | 58.5      | 1.5   |
| <u>Artemisia ludoviciana</u>       |                 |           |       | 94.5      | 1.9   | 430.0     | 10.8  |
| <u>Carex spp.</u>                  |                 | 10.0      | 0.2   | 124.0     | 2.5   | 33.0      | 0.8   |
| <u>Cirsium arvense</u>             |                 | 94.5      | 1.7   |           |       |           |       |
| <u>Rosa woodsii</u>                |                 |           |       | 578.5     | 11.5  |           |       |
| <u>Symphoricarpos occidentalis</u> |                 | 2871.5    | 50.3  | 2775.5    | 55.2  | 2821.5    | 70.8  |
| Unclassified forbs                 |                 | 5.0       | 0.1   |           |       | 45.0      | 1.1   |
| Total                              |                 | 5713.0    | 100.1 | 5024.0    | 100.0 | 3985.5    | 100.0 |

Table 10. Species Composition and Amount of Vegetation at Channel 6-69, 2-69, 5-69, and 8<sup>a</sup>-69 Nest Sites, Gorham Study Area, North Dakota. Vegetation Sampled by Clipped Hoop Method, Spring 1970.

| Species                            | Channel<br>Date | 6 <sup>c</sup><br>5-6-70 |       | 2<br>5-16-70 |       | 5<br>5-16-70 |       | 8 <sup>a</sup><br>5-18-70 |       |
|------------------------------------|-----------------|--------------------------|-------|--------------|-------|--------------|-------|---------------------------|-------|
|                                    |                 | lbs./acre                | %     | lbs./acre    | %     | lbs./acre    | %     | lbs./acre                 | %     |
| <u>Agropyron cristatum</u>         |                 |                          |       | 1814.5       | 100.0 | 2804.5       | 100.0 |                           |       |
| <u>Agropyron smithii</u>           |                 | 787.5                    | 48.9  |              |       |              |       | 4980.5                    | 100.0 |
| <u>Bromus inermis</u>              |                 |                          |       |              |       |              |       |                           |       |
| <u>Rosa woodsii</u>                |                 | 96.0                     | 6.0   |              |       |              |       |                           |       |
| <u>Symphoricarpos occidentalis</u> |                 | 725.5                    | 45.0  |              |       |              |       |                           |       |
| Unclassified forbs                 |                 | 2.5                      | 0.1   |              |       |              |       |                           |       |
| Total                              |                 | 1611.5                   | 100.0 | 1814.5       | 100.0 | 2804.5       | 100.0 | 4980.5                    | 100.0 |

<sup>a</sup>Initial nest

<sup>c</sup>Located at the base of a Sheperdia argentea bush

Table 11. Species Composition and Amount of Vegetation at Channel 1<sup>a</sup>-70, 1<sup>b</sup>-70 and 2-70 Nest Sites, Golden Study Area, North Dakota. Vegetation Sampled by Clipped Hoop Method, Summer 1970.

| Species                            | Channel<br>Date | 1 <sup>a</sup> c<br>6-9 |      | 1 <sup>b</sup><br>7-18 |       | 2<br>6-26 |       |
|------------------------------------|-----------------|-------------------------|------|------------------------|-------|-----------|-------|
|                                    |                 | lbs./acre               | %    | lbs./acre              | %     | lbs./acre | %     |
| <u>Agropyron cristatum</u>         |                 |                         |      | 2862.5                 | 68.0  |           |       |
| <u>Agropyron repens</u>            |                 |                         |      |                        |       | 1500.0    | 17.7  |
| <u>Agropyron smithii</u>           |                 |                         |      | 186.0                  | 4.4   |           |       |
| <u>Boutelona gracilis</u>          |                 |                         |      |                        |       | 12.0      | 0.1   |
| <u>Poa palustris</u>               |                 |                         |      | 68.0                   | 1.6   |           |       |
| <u>Stipa viridula</u>              |                 | 731.0                   | 17.3 | 75.0                   | 1.8   |           |       |
| <u>Achillea lanulosa</u>           |                 |                         |      | 221.0                  | 5.3   |           |       |
| <u>Artemisia ericardes</u>         |                 | 1.5                     | tr.  |                        |       |           |       |
| <u>Artemisia frigida</u>           |                 |                         |      | 157.0                  | 3.7   |           |       |
| <u>Artemisia ludoviciana</u>       |                 | 201.5                   | 4.8  |                        |       |           |       |
| <u>Carex spp.</u>                  |                 | 10.5                    | 0.2  |                        |       |           |       |
| <u>Lactuca pulchella</u>           |                 | 5.5                     | 0.1  |                        |       |           |       |
| <u>Medicago sativa</u>             |                 |                         |      |                        |       | 6716.0    | 79.4  |
| <u>Psoralea argophylla</u>         |                 |                         |      | 631.5                  | 15.0  |           |       |
| <u>Rosa arkansana</u>              |                 |                         |      |                        |       | 218.0     | 2.6   |
| <u>Rosa woodsii</u>                |                 | 229.5                   | 5.4  |                        |       |           |       |
| <u>Symphoricarpos occidentalis</u> |                 | 3022.5                  | 71.5 |                        |       |           |       |
| <u>Unclassified forbs</u>          |                 | 26.5                    | 0.6  | 7.5                    | 0.2   | 17.0      | 0.2   |
| Total                              |                 | 4228.5                  | 99.9 | 4208.5                 | 100.0 | 4863.0    | 100.0 |

<sup>a</sup>Initial nest

<sup>c</sup>Located at the base of Sheperdia argentea bush

<sup>b</sup>Renest

Table 12. Species Composition and Amount of Vegetation at Channel 3<sup>a</sup> - 70, 3<sup>b</sup> - 70, and 6-70 Nest Sites, Gorham Study Area, North Dakota. Vegetation Sampled by Clipped Hoop Method, Summer 1970.

| Species                            | Channel<br>Date | 3 <sup>a</sup><br>6-9 |      | 3 <sup>b</sup><br>7-17 |      | 6<br>6-23 |      |
|------------------------------------|-----------------|-----------------------|------|------------------------|------|-----------|------|
|                                    |                 | lbs./acre             | %    | lbs./acre              | %    | lbs./acre | %    |
| <u>Agropyron cristatum</u>         |                 |                       |      |                        |      | 2903.5    | 46.3 |
| <u>Agropyron repens</u>            |                 | 3073.5                | 60.6 | 2226.5                 | 29.0 |           |      |
| <u>Agropyron trachycaulum</u>      |                 |                       |      | 1409.5                 | 18.4 | 18.0      | 0.3  |
| <u>Avena fatua</u>                 |                 | 6.0                   | 0.1  | 25.0                   | 0.3  |           |      |
| <u>Bromus inermis</u>              |                 |                       |      |                        |      | 3179.0    | 50.7 |
| <u>Festuca octoflora</u>           |                 |                       |      |                        |      | 51.5      | 0.8  |
| <u>Muhlenbergia richardsonis</u>   |                 |                       |      | 111.0                  | 1.4  | 90.5      | 1.4  |
| Annual weeds                       |                 | 1949.0                | 38.4 |                        |      |           |      |
| <u>Rosa woodsii</u>                |                 |                       |      | 1293.5                 | 16.9 |           |      |
| <u>Symphoricarpos occidentalis</u> |                 |                       |      | 2498.0                 | 32.5 |           |      |
| Unclassified forbs                 |                 | 41.5                  | 0.8  | 110.5                  | 1.4  | 23.0      | 0.4  |
| Total                              |                 | 5070.0                | 99.9 | 7674.0                 | 99.9 | 6265.5    | 99.9 |

<sup>a</sup>Initial nest

<sup>b</sup>Renest

Table 13. Species Composition and Amount of Vegetation at Channel 10<sup>a</sup>-70, 10(nest bowl) - 70, and 10<sup>b</sup>-70 Nest Sites, Gorham Study Area, North Dakota. Vegetation Sampled by Clipped Hoop Method, Summer 1970.

| Species                    | Channel<br>Date | 10 <sup>a</sup> |       | 10 <sup>c</sup> |       | 10 <sup>b</sup> |       |
|----------------------------|-----------------|-----------------|-------|-----------------|-------|-----------------|-------|
|                            |                 | 6-18            |       | 6-29            |       | 7-17            |       |
|                            |                 | lbs./acre       | %     | lbs./acre       | %     | lbs./acre       | %     |
| <u>Agropyron cristatum</u> |                 |                 |       | 411.5           | 12.0  | 332.5           | 9.9   |
| <u>Agropyron smithii</u>   |                 | 223.5           | 5.2   | 1125.5          | 33.0  | 699.5           | 20.9  |
| <u>Bromus japonicus</u>    |                 | 111.5           | 2.6   |                 |       |                 |       |
| <u>Koeleria cristata</u>   |                 |                 |       | 381.0           | 11.2  | 39.5            | 1.2   |
| <u>Poa arida</u>           |                 | 542.5           | 12.7  |                 |       |                 |       |
| <u>Poa glauicifolia</u>    |                 |                 |       |                 |       | 1364.5          | 40.8  |
| <u>Poa pratensis</u>       |                 |                 |       | 105.5           | 3.1   |                 |       |
| <u>Stipa viridula</u>      |                 | 3300.0          | 77.1  | 1019.5          | 29.9  |                 |       |
| Annual weeds               |                 | 29.0            | 0.7   |                 |       | 519.5           | 15.5  |
| <u>Carex spp.</u>          |                 | 21.0            | 0.5   |                 |       |                 |       |
| <u>Psoralea argophylla</u> |                 |                 |       |                 |       | 130.5           | 3.9   |
| Unclassified forbs         |                 | 52.0            | 1.2   | 89.0            | 2.6   | 260.5           | 7.8   |
| <u>Vicia americana</u>     |                 |                 |       | 283.0           | 8.3   |                 |       |
| Total                      |                 | 4279.5          | 100.0 | 3415.0          | 100.1 | 3346.5          | 100.0 |

<sup>a</sup>Initial nest

<sup>b</sup>Renest

<sup>c</sup>Nest bowl

Table 14. Species Composition and Amount of Vegetation at Channel 8-70, 12<sup>a</sup>-70, and 10<sup>i</sup>-70 Nest Sites, Gorham Study Area, North Dakota. Vegetation Sampled by Clipped Hoop Method, Summer 1970.

| Species                     | Channel<br>Date |  | 8<br>6-22 |       | 12 <sup>a</sup><br>6-18 |       | 12 <sup>b</sup><br>7-27 |      | 10 <sup>i</sup> c<br>6-26 |       |
|-----------------------------|-----------------|--|-----------|-------|-------------------------|-------|-------------------------|------|---------------------------|-------|
|                             |                 |  | lbs./acre | %     | lbs./acre               | %     | lbs./acre               | %    | lbs./acre                 | %     |
| Agropyron cristatum         |                 |  | 6510.5    | 96.0  | 4941.5                  | 95.2  | 4724.0                  | 88.7 |                           |       |
| Agropyron smithii           |                 |  |           |       | 30.0                    | 0.6   |                         |      | 465.0                     | 6.2   |
| Koeleria cristata           |                 |  |           |       |                         |       | 160.5                   | 3.0  |                           |       |
| Panicum virgatum            |                 |  |           |       |                         |       |                         |      | 25.5                      | 0.3   |
| Panicum wilcoxianum         |                 |  |           |       |                         |       |                         |      | 6.5                       | 0.1   |
| Stipa comata                |                 |  |           |       |                         |       | 350.0                   | 6.6  |                           |       |
| Ambrosia psilostachya       |                 |  |           |       | 27.0                    | 0.5   |                         |      |                           |       |
| Artemisia frigida           |                 |  | 3.5       | 0.1   | 139.0                   | 2.7   |                         |      |                           |       |
| Artemisia ludoviciana       |                 |  |           |       |                         |       |                         |      | 191.5                     | 2.6   |
| Carex spp.                  |                 |  |           |       | 8.5                     | 0.2   |                         |      | 2.0                       | tr.   |
| Psoralea argophylla         |                 |  |           |       | 11.5                    | 0.2   |                         |      |                           |       |
| Rosa arkansana              |                 |  | 270.5     | 4.0   |                         |       |                         |      |                           |       |
| Selaginella densa           |                 |  |           |       | 25.5                    | 0.5   |                         |      |                           |       |
| Symphoricarpos occidentalis |                 |  |           |       |                         |       |                         |      | 6806.5                    | 90.7  |
| Unclassified forbs          |                 |  |           |       | 8.5                     | 0.2   | 88.5                    | 1.6  | 1.6                       | 0.1   |
| Total                       |                 |  | 6784.5    | 100.1 | 5191.5                  | 100.1 | 5323.0                  | 99.9 | 7501.0                    | 100.0 |

<sup>a</sup>Initial nest

<sup>b</sup>Renest

<sup>i</sup>Intermittent signal channel

<sup>c</sup>Located at the base of a Sheperdia argentea bush

Table 15. Characteristics of Cover at Sharp-tailed Grouse Nest Sites, Gorham Study Area, North Dakota,  
Based on 2 X 1 Foot Cover Board, 1969-1970.

| Site               | Density in<br>percent | Height in<br>inches | Dominants   | Proximity<br>to bushy<br>draw<br>(yards) | Percent<br>Slope | Grazing             |
|--------------------|-----------------------|---------------------|---|--|------------------|---------------------|
| 1-69               | 100<br>20             | 4<br>14             | <u>Symphoricarpos occidentalis</u><br><u>Stipa viridula</u> | 0  | 10               | Summer              |
| 2-69               | 100<br>40             | 3<br>12             | <u>Agropyron cristatum</u>                                  | 150                                      | 0                | Summer              |
| 3-69               | 100<br>15             | 4<br>15             | <u>Symphoricarpos occidentalis</u><br><u>Stipa viridula</u> | 0  | 20               | Summer              |
| 4 <sup>a</sup> -69 | 100<br>40             | 3<br>12             | <u>Agropyron cristatum</u><br><u>Stipa comata</u>           | 150                                      | 5                | Summer              |
| 4 <sup>b</sup> -69 | 100<br>90             | 10<br>20            | <u>Melilotus officinalis</u>                                | 475                                      | 0                | Hay-land            |
| 5-69               | 100<br>40             | 3<br>12             | <u>Agropyron cristatum</u>                                  | 75                                       | 0                | Summer              |
| 6-69               | 100<br>20             | 4<br>43             | <u>Sheperdia argentea</u><br><u>Agropyron smithii</u>       | 0  | 45               | Summer              |
| 7-69               | 100<br>30             | 3<br>15             | <u>Symphoricarpos occidentalis</u><br><u>Stipa viridula</u> | 0  | 20               | Summer              |
| 8 <sup>a</sup> -69 | 100<br>40             | 7<br>14             | <u>Bromus inermis</u>                                       | 330                                      | 5                | Winter <sup>d</sup> |
| 8 <sup>b</sup> -69 | 100<br>80             | 4<br>24             | <u>Melilotus officinalis</u><br><u>Bromus inermis</u>       | 800                                      | 10               | Hay-land            |



Table 15 Continued.

| Site                | Density in percent | Height in inches | Dominants   | Proximity to bushy draw (yards) | Percent Slope | Grazing             |
|---------------------|--------------------|------------------|---|---------------------------------|---------------|---------------------|
| 10 <sup>a</sup> -69 | 100<br>70          | 4<br>15          | <u>Symphoricarpos occidentalis</u>                              | 0                               | 20            | Winter <sup>d</sup> |
| 10 <sup>b</sup> -69 | 100<br>30          | 3<br>14          | <u>Symphoricarpos occidentalis</u><br><u>Stipa viridula</u>     | 100                             | 5             | Winter <sup>d</sup> |
| 1 <sup>a</sup> -70  | 100<br>40          | 3<br>42          | <u>Sheperdia argentea</u><br><u>Symphoricarpos occidentalis</u> | 0                               | 25            | Summer              |
| 1 <sup>b</sup> -70  | 100<br>50          | 3<br>18          | <u>Agropyron cristatum</u><br><u>Agropyron smithii</u>          | 45                              | 5             | Summer              |
| 2-70                | 100<br>30          | 6<br>16          | <u>Medicago sativa</u><br><u>Agropyron repens</u>               | 350                             | 15            | Hay-land            |
| 3 <sup>a</sup> -70  | 100<br>70          | 10<br>28         | <u>Agropyron repens</u><br>Annual weeds                         | 300                             | 10            | Non-use             |
| 3 <sup>b</sup> -70  | 100<br>70          | 9<br>28          | <u>Symphoricarpos occidentalis</u><br><u>Agropyron repens</u>   | 130                             | 0             | Non-use             |
| 6-70                | 100<br>80          | 15<br>30         | <u>Bromus inermis</u><br><u>Agropyron cristatum</u>             | 1760                            | 15            | Non-use             |
| 8-70                | 100<br>60          | 6<br>25          | <u>Agropyron cristatum</u><br><u>Rosa arkansana</u>             | 300                             | 20            | Summer              |
| 10 <sup>a</sup> -70 | 100<br>40          | 5<br>12          | <u>Stipa viridula</u><br><u>Poa arida</u>                       | 120                             | 10            | Summer              |
| 10 <sup>c</sup> -70 | 100<br>15          | 5<br>16          | <u>Agropyron cristatum</u><br><u>Stipa viridula</u>             | 120                             | 5             | Summer              |

Table 15 Continued.

| Site                | Density in percent | Height in inches | Dominants                          | Proximity to bushy draw (yards) | Percent Slope | Crazing             |
|---------------------|--------------------|------------------|------------------------------------|---------------------------------|---------------|---------------------|
| 10 <sup>b</sup> -70 | 100                | 4                | <u>Agropyron cristatum</u>         | 130                             | 5             | Summer              |
|                     | 25                 | 21               | <u>Poa glaucifolia</u>             |                                 |               |                     |
| 12 <sup>a</sup> -70 | 100                | 2                | <u>Agropyron cristatum</u>         | 220                             | 5             | Summer              |
|                     | 60                 | 16               | <u>Agropyron smithii</u>           |                                 |               |                     |
| 12 <sup>b</sup> -70 | 100                | 4                | <u>Agropyron cristatum</u>         | 210                             | 20            | Summer              |
|                     | 50                 | 24               | <u>Stipa comata</u>                |                                 |               |                     |
| 10 <sup>i</sup> -70 | 100                | 4                | <u>Sheperdia argentea</u>          | 0                               | 15            | Summer <sup>d</sup> |
|                     | 60                 | 70               | <u>Symphoricarpos occidentalis</u> |                                 |               |                     |

<sup>a</sup>Initial nest<sup>b</sup>Renest<sup>c</sup>Nest bowl<sup>d</sup>Lightly grazed; portions also hayed<sup>i</sup>Intermittent signal channel

## Brood movement and cover selection

Cover characteristics and adjacent land-use data for 37 brood locations observed during the summers of 1969 and 1970 are presented in Table 16. These observations were made only in open areas serviced by roads or trails.

Eight radio equipped brooding hens were followed for periods of 4 to 61 days. Channel 2-69, which hatched eight of fifteen eggs from her nest in a crested wheatgrass field on June 13, remained within 50 yards of the nest during the following day. On June 15 the brood moved 0.3 mi. south southwest in the crested wheatgrass field and remained there for two days. On June 17 they moved 0.4 mi. north northeast, past the nest site, into a cultivated field grown over with sweet clover and frenchweed (*Thlaspi arvense*). For the next three days the brood used this field and adjacent wooded and brushy draws. This brood moved 0.5 mi. south on June 21 to an area of sweet clover and brush within the crested wheatgrass field. After three days they moved 0.3 mi. southeast to an adjacent crested wheatgrass field on June 24. During the next three days a continuous rain apparently resulted in the loss of the brood. The hen moved 0.5 mi. further south remaining in a large pasture with deep draws for the remainder of the summer and was collected at this location on September 20 (Figure 3).

Channel 8-69, in her reneesting success, hatched all nine eggs from a nest under a sweet clover bush in a smooth brome-alfalfa field July 17. This brood moved 200 yards north to the edge of an alfalfa field through July 19 and was found approximately 100 yards east of the nest site in an adjacent smooth brome-alfalfa field on July 20. The next day they had moved 0.4 mi. northwest to an alfalfa field where windrowing operations apparently resulted in the loss of the entire brood, because they were never seen with the hen again. The female spent the remainder of the summer in the draw area in which both the initial and reneest had been located (Figure 4).

On July 20, channel 10-69 hatched six of eleven eggs from a nest located on a buckbrush flat. These were the result of an attempt to reneest. She immediately moved the brood 100 yards north to a wheat (*Triticum* spp.) field and remained there until July 31. During the first four days of August the brood remained within 50 yards of the nest site in a pasture and used prairie thistle (*Cirsium undulatum*) and sweet clover for cover. The brood then moved 0.4 mi. southwest in this pasture, locating in crested wheatgrass, sweet clover, alfalfa, and white sage (*Artemisia ludoviciana*) patches from August 5-8. From August 9-18 they remained within 100 yards of the nest site and utilized prairie thistle and sweet clover cover. The hen and brood relocated in this pasture along a buckbrush draw 0.4 mi. south of the nest site on August 19-20. The study was terminated on August 20 and the hen was collected at the same location on September 20 (Figure 5).

Channel 1-70 hatched seven of ten eggs from her reneest between two crested wheatgrass bunches July 15. After an initial move to a crested wheatgrass hilltop 0.4 mi. south of the nest site, this brood remained in the vicinities of the hilltop and the nest site through July 31 with occasional movements up to 0.5 mi. between these sites. August 1 and 2 were spent near a draw 0.5 mi. north of the nest site. The brood moved 0.7 mi. northeast to a native hayland on August 3 and remained there until the hen was taken by an avian predator on August 11. Throughout the first week this brood utilized crested wheatgrass exclusively for cover; during the second, third, and fourth weeks more dense vegetation (buckbrush and sweet clover) was used, particularly during the heat of the day. Throughout the period the brood roosted on upland sites (Figure 6).

Table 16. Cover Characteristics and Adjacent Land-use at 37 Sharp-tailed Grouse Brood Locations, Gorham Study Area, Southwestern North Dakota, 1969-1970.

| Type       | No. of broods | Cover height |        |          | Density |        |       | Species Composition |        |       | Adjacent Land-use |               |                         |                             |                     |              |
|------------|---------------|--------------|--------|----------|---------|--------|-------|---------------------|--------|-------|-------------------|---------------|-------------------------|-----------------------------|---------------------|--------------|
|            |               | Under 12"    | 12-30" | Over 30" | Sparse  | Medium | Heavy | Native              | Exotic | Mixed | Summer grazed     | Winter grazed | Winter grazed and hayed | Non-use with portions hayed | Non-use Small grain | Summerfallow |
| Grassland  | 20            | 9            | 10     | 1        | 3       | 12     | 5     | 12                  | 2      | 6     | 13                | 2             | 2                       | 1                           | 2                   |              |
| Brush      | 6             |              | 4      | 2        |         | 2      | 4     | 5                   | 1      |       | 4                 |               |                         |                             | 2                   |              |
| Roadside   | 6             | 1            | 3      | 2        |         | 2      | 4     | 1                   | 4      | 1     | 1                 |               |                         | 1                           | 3                   | 1            |
| Cultivated | 2             | 1            | 1      |          |         | 1      | 1     |                     | 2      |       |                   |               |                         | 2                           |                     |              |
| Hayland    | 3             | —            | 1      | 2        | —       | —      | 3     | —                   | 3      | —     | —                 | —             | 2                       | —                           | —                   | 1            |
| Total      | 37            | 11           | 19     | 7        | 3       | 17     | 17    | 18                  | 12     | 7     | 18                | 2             | 4                       | 4                           | 7                   | 1 1          |




Figure 3. Channel 2-69 Brood Movement, Gorham Study Area North Dakota, 1969.




Figure 4. Channel 8-69 Brood Movement, Gorham Study Area, North Dakota, 1969.




Figure 5. Channel 10-69 Brood Movement, Gorham Study Area, North Dakota, 1969.




Figure 6. Channel 1-70 Brood Movement, Gorham Study Area, North Dakota, 1970.



Channel 2-70 hatched 13 of 15 eggs in an alfalfa field on June 25 and moved her brood 0.3 mi. north to an over-grazed pasture during the following week. Upland sites were used for cover during this period. On July 3 the brood moved 0.8 mi. east to a wooded draw and remained there except for short daily movements until August 3. Contact with this brood was lost until August 12 when it was relocated in a wooded draw 1.5 mi. northeast of the original draw; its activities centered here until the termination of the study on August 25. From July 3 on, its activities were centered in wooded draws except for feeding during cool portions of the day and roosting at night (Figure 7).

On June 22 channel 6-70 hatched her entire clutch of 15 eggs. The next five days were spent along a grassy ridge 0.2 mi. north of the nest site. On July 28 the brood was moved 0.8 mi. north to a heavily grazed pasture and two days later 0.5 mi. west in this pasture where brushy vegetation was occasionally utilized for cover. From July 2 through July 7 the brood moved 1.2 mi. north-east into the Fairfield Common Pasture where the hen was lost to an avian predator July 8 (Figure 8).

Channel 8-70 hatched 13 eggs on June 22. The following two days were spent in the crested wheatgrass within 0.2 mi. of the nest site. On June 25 the brood moved 0.3 mi. south to an adjacent crested wheatgrass field and the next day moved an additional 0.4 mi. south in the same field. On July 1 the brood traveled 1.0 mi. southwest to a native pasture where it remained through July 2. The following day the brood was located 0.7 mi. north in sweet clover cover adjacent to a wooded draw. Contact with the brood was lost until July 6 when the brood was relocated in a wooded draw only 0.4 mi. southwest of the nest site. The brood remained here until July 9 when it again moved into the adjacent crested wheatgrass field, south of the nest site. Following a week in this area, during which brushy vegetation was used only during hot weather, the brood traveled 1.0 mi. northeast on July 16 to a native hayland with considerable sweet clover. This cover was utilized until July 23 when the brood moved 1.5 mi. south to a wooded draw which was its center of activity until the hen was killed by a mammalian predator on August 8 (Figure 9).

Channel 10<sup>1</sup>-70 hatched 14 of 15 eggs from a nest located in a buffalo berry-buckbrush site. After one day spent in the vicinity of the adjacent draw the brood moved 0.6 mi. southwest to a crested wheatgrass field July 1 only to return to the native pasture July 3. The following week was spent here and at the edge of the adjacent Fairfield Common Pasture where sweet clover was the principal cover utilized. From July 11 until the hen was taken by a mammalian predator July 24 the brood remained 0.4 mi. north in the native hayland also used by channel 8-70 and utilized sweet clover as the primary cover (Figure 10).

Daily movements of these eight sharptail broods were generally within 1.0 mi. of their respective nest sites. Occasional daily movements of up to 1.0 mi. were noted in the study; broods less than a week old moved up to 0.8 mi. Table 17 summarizes the brood monitoring data, including home ranges.

Development of flight capacity at an early stage was observed in two broods. Broods of channel 6-70 and 8-70 were able to fly up to 30 yards at eight and nine days of age, respectively.




Figure 7. Channel 2-70 Brood Movement, Gorham Study Area, North Dakota, 1970.




Figure 8. Channel 6-70 Brood Movement, Gorham Study Area, North Dakota, 1970.




Figure 9. Channel 8-70 Brood Movement, Gorham Study Area, North Dakota, 1970.




Figure 10. Channel 10<sup>i</sup>-70 Brood Movement, Gorham Study Area, North Dakota, 1970.

Table 17. Home Range<sup>a</sup> and Daily Movements<sup>b</sup> of Eight Sharp-tailed Grouse Broods, Gorham Study Area, North Dakota, 1969-1970.

| Weeks                         | Broods |      |       |      |      |      |      |                     |
|-------------------------------|--------|------|-------|------|------|------|------|---------------------|
|                               | 2-69   | 8-69 | 10-69 | 1-70 | 2-70 | 6-70 | 8-70 | 10 <sup>i</sup> -70 |
| Acres                         |        |      |       |      |      |      |      |                     |
| 1                             | 50     | 80   | 20    | 60   | 20   | 50   | 45   | 50                  |
| 2                             | 120    | -    | 20    | 50   | 40   | 60   | 100  | 40                  |
| 3                             | -      | -    | 40    | 80   | 60   | 120  | 110  | 35                  |
| 4                             | -      | -    | 100   | 40   | 60   | -    | 130  | 45                  |
| 5                             | -      | -    | -     | -    | 80   | -    | 120  | -                   |
| 6                             | -      | -    | -     | -    | 120  | -    | 60   | -                   |
| 7                             | -      | -    | -     | -    | 120  | -    | 70   | -                   |
| 8                             | -      | -    | -     | -    | 60   | -    | ---  | -                   |
| 9                             | -      | -    | -     | -    | 70   | -    | ---  | -                   |
| Miles                         |        |      |       |      |      |      |      |                     |
| 1                             | 0.4    | 0.4  | 0.1   | 0.4  | 0.3  | 0.8  | 0.4  | 0.6                 |
| 2                             | 0.5    | -    | 0.1   | 0.4  | 0.8  | 0.5  | 1.0  | 0.6                 |
| 3                             | -      | -    | 0.4   | 0.4  | 0.4  | 0.3  | 0.4  | 0.4                 |
| 4                             | -      | -    | 0.4   | 0.7  | 0.2  | -    | 1.0  | 0.2                 |
| 5                             | -      | -    | -     | -    | 0.4  | -    | 0.8  | -                   |
| 6                             | -      | -    | -     | -    | 0.3  | -    | 0.3  | -                   |
| 7                             | -      | -    | -     | -    | 0.5  | -    | 0.4  | -                   |
| 8                             | -      | -    | -     | -    | 0.2  | -    | ---  | -                   |
| 9                             | -      | -    | -     | -    | 0.2  | -    | ---  | -                   |
| Total<br>acre-<br>age<br>used | 120    | 80   | 150   | 200  | 300  | 200  | 500  | 160                 |

<sup>a</sup>In acres

<sup>b</sup>In miles

<sup>i</sup>Intermittent signal channel



Three brood hens, channel 2-70, 8-70, and 10<sup>i</sup>-70, were monitored continuously for two separate days to determine times of activities and cover used. A generalized summary of the daily activities of all broods observed in the study is presented in Figure 11. All broods had similar daily activity patterns, which began with intensive feeding immediately after sunrise in open areas. This was followed by a period of intermittent resting and feeding in increasingly dense cover. By mid-day the brood was resting in dense cover in draws. Subsequently, this was followed by intermittent feeding and resting in decreasingly dense cover which phases into an intense evening feeding period in light cover. Roosting was confined to upland sites in light cover. Wind, precipitation, and variations in daily temperatures affected the normal activities.




Figure 11. Daily Activities and Cover Selection of Sharp-tailed Grouse Broods, Gorham Study Area, North Dakota, 1969-1970.

# DISCUSSION

## Trapping

Various techniques were used to recapture instrumented hens for battery replacement. One brooding hen was netted with a mobile cannon-net. Of 13 incubating hens recaptured on their nests, one was caught by hand, seven by hoop net, and five by bow-net trap. Bow-net problems include malfunction in dense vegetation and the interval required to allow the hen to return after flushing. The problems involved with hoop netting include the difficulty of approaching wary birds and netting them quickly and properly in order to prevent injury to the bird or nest. In an effort to create as little disturbance as possible at the nest site and to limit the number of trails into the nest, hoop net capture was attempted only once. If this was unsuccessful, the nest trap was used to assure that only one more disturbance would take place. With one exception, all captures were carried out towards the end of incubation to prevent desertion. One hen deserted after her non-functional transmitter was replaced early in incubation.

Active dancing had been observed on ground 5, a cultivated site, during the springs of 1966-1968. However, active dancing concentrations could not be located here in 1969 although small groupings of birds were occasionally seen. In 1970, a new ground was found one-half mi. to the west at a corner of a heavily grazed pasture. It was assumed that this was a relocation of ground 5 from the former cultivated site, because one male taken here had been previously banded at the cultivated site during the spring of 1967.

## Telemetry

Telemetry equipment did not cause observable unusual behavioral effects, and normal nesting, incubating, and hatching or reneesting took place. A copulation between a radioed hen and a male on a dancing ground only two days after the hen's release indicated a rapid adjustment to the radio package. Previous studies have reported that instrumentation of game birds does not effect normal reproductive activity (Brown, 1965; Cebula, 1966; Kuck, 1966; Bernhoff, 1969).

## Hen dispersal and nesting

One hen may have attended two different dancing grounds during the spring of 1969. Channel 3-69 had been trapped and released April 30 on dancing ground 4. She remained for 20 days in an area 1.4 mi. to the northeast, in the vicinity of a previously undetected dancing ground. On May 20 the hen moved 0.6 mi. further northeast to a brushy draw in which her nest was found May 30. Because the new dancing ground was not located until May 21, ch. 3-69's attendance on this ground was not determined; however, the hen was frequently found in association with other grouse from this dancing ground area.

During 1969, a year with average weather conditions, hen attendance at dancing grounds was approximately three weeks. Thus, the nesting was extended; initial attempts were made throughout the period from April 25 through May 28. Therefore, in 1969 the period of nesting was considered to be of normal duration. In 1970, however, 26 inches of snow were recorded on the study area between April 10 and May 4. This apparently curtailed dancing ground activity which resulted in a short peak of attendance by hens and a subsequent concentration of all but one of eight initial nesting attempts into one four day period between May 12 and May 16. Predation may have extended the period in which nesting was

attempted and thus reduced the possibility that all nests and broods would be eliminated by the same general disaster. This is in accord with the findings of Cartwright (1944) and Symington and Harper (1957) that predation may be a mechanism which operates in prolonging the period when nesting is attempted.

Hamerstrom (1939), Cartwright (1944), Ammann (1957), and Blus and Walker (1966) reported that the reproductive success of prairie grouse depends upon initial nesting efforts because they do not persistently renest, however, considerable renesting was observed in the present study. Seven of nine birds whose nests were destroyed by predators, renested. The two other birds apparently did not renest, however their nests were found in early June and may have been the product of renesting.

Renesting was more successful than initial nesting. Four of six renests (not including a desertion caused by early transmitter replacement) hatched while only 7 of 17 initial nests hatched. As the season progressed the amount of available nesting cover vastly increased. Therefore, if the numbers of predators in the area remained constant, the additional cover probably increased renesting success.

#### Vegetation at nest sites

Of 25 nest sites (including ch. 10-70's nest bowl), ten were associated with crested wheatgrass, five with buckbrush, three with buffalo berry, two with sweet clover, two with quackgrass, and one each with smooth brome, alfalfa, and green needlegrass. The only comparable characteristic found at all sites was the height of the cover. Vegetation at all sites, other than those associated with buckbrush and buffalo berry was at least 12 inches tall over a uniform area. Nest sites associated with buckbrush and buffalo berry were at locations along edges where the height of the vegetation was from 14 to 70 inches. Therefore, it would seem that sharptail hens require uniform vegetation not less than 12 inches or patchy vegetation not less than 14 inches in height for nesting. The predominance of heavily grazed pasture on the Gorham Study Area where the height of vegetation was considerably less than a uniform 12 inches may have forced many hens to seek taller patchy vegetation for nesting sites. In most cases these were brushy draws which are definite predator lanes. This was reflected in the destruction of 5 of 7 nests located in brushy draws. In addition, only 2 of 11 clutches which were successfully hatched were from nests located near these draws.

Crested wheatgrass was the dominant species present at both initial and reneest sites. This may indicate that this was preferred nesting cover because sharptails tend to locate nests in crested wheatgrass. On the other hand wheatgrass may be ideal initial nesting cover because it over-winters well and, thus, provides residual cover. Residual vegetation comprised between 10 to 90 percent of nesting cover at initial nests. In addition, six of eight nests associated with crested wheatgrass hatched. Conversely buckbrush and buffalo berry may be considered as less preferred nesting cover and of questionable value. Although 8 of 25 nests were associated with buckbrush and/or buffalo berry only two hatched.

## Predation

Predation apparently plays an important role in limiting sharptail reproduction. Twelve of 24 nests located in the study were lost to predation. The average length of time between the last visual observation of the nest and predation was approximately eight days with a range from 3 to 15 days; hens were monitored daily. In many cases a rain shower or thunderstorm intervened between this observation and predation. It seems unlikely that predators could trail the investigator to the nest sites under such conditions, however, Brown (1966b) found that increased nesting hen predation by coyotes was apparently caused by trailing. In two, possibly three, of the 12 nest predations the hen was taken. This was between ten and twenty percent of the 17 instrumented hens.

Brooding hens are apparently more susceptible to predation than non-brooding hens. Four of eight hens with broods were taken by predators, while only one of five non-brooding hens was preyed upon. This may be due to the fact that brooding hens tend to hold tighter when approached and can be readily taken by the predator.

## Brood movement and cover selection

Of 11 hatching broods only portions of two (three young) were with the hens at the termination of the study in late August. Severe weather apparently caused the loss of three broods. Channel 2-69's chicks were only 11 days old when a cold three day rain occurred; they were never again seen. Channel 6-69 and 12-70 were in the process of hatching during severe thunderstorms with high winds, heavy rain, and a large amount of small hail. Both nests were observed within 12 hours after the storms. A few hatched young had been washed down the slope and the remainder of the broods had only partially emerged from the shells. Haying operations resulted in the loss of one brood, channel 8-69's, which was located in an alfalfa field at the time it was being windrowed. The hen was flushed from the field and one of eight chicks was found alive in a windrow; the hen was never seen with the brood again. One brood died in the early morning cold on the day following hatching during a prolonged attempt to capture the hen. The hens of the other four broods (28 young) were lost to predation (two mammalian and two avian). It was not determined if any or all of these young lived until the study ended although the broods ranged from 16 to 47 days of age at the time of predation and most were increasingly independent of the hens.

Young broods less than two weeks old apparently were dependent only on the hens for shelter as they generally moved to and remained in closely grazed pastures of short sparse vegetation. Subsequently, there was an increasing dependence on more dense cover because the broods moved to areas which furnished available protective cover, usually brushy or wooded draws.

The daily movements and ranges varied throughout the study but a general pattern was followed by most broods. At an early age the broods moved relatively short distances and had small ranges; at three to four weeks of age brood movements and ranges reached maximums. Although only two broods were monitored at ages greater than five weeks, both localized in close proximity to draws and, therefore, had reduced daily movements and ranges.

#### RECOMMENDATIONS

If the activities of this small group of radio-equipped hens are indicative of the true situation, then one form of improved land management can be suggested; this would also be effective grouse management. Sharptail populations might be substantially increased by regulating grazing to provide uniform areas with 12 inch high vegetation for nesting cover. This can be provided either by establishing small exclosures in heavily grazed areas or by regulating grazing rotation so that it will provide suitable spring nesting habitat each year.

The following recommendations are suggested:

- (1) Continue the investigation to determine whether an individual female attends more than one dancing ground and to investigate the possible effect this will have on the location of the initial and subsequent nesting.
- (2) Examine the allotment records of the last five years, for all public lands included in the study area, to determine if there is a correlation between grazing pressure and a concentration of grouse nesting.
- (3) Examine existing cattle exclosures in grouse habitat in southwestern North Dakota to determine whether increased nesting activity and greater hatching success are realized on such areas.
- (4) Establish cattle exclosures to determine whether they will enhance sharptail nesting on the study area. This data could be readily compared with data concerning nesting which has been collected over a period of four years.
- (5) Determine the condition and trend of the vegetation and soil of the area in the vicinity of all nest sites by utilizing paced line transects.
- (6) Determine the time the female spends on the nest in laying and incubating the clutch and monitor temperature and humidity in sharptail nests with battery operated timing devices and hygrothermographs.
- (7) Record the density of vegetation, the precise time of day and the temperature at each brood location to determine whether there is a correlation between these factors.
- (8) Secure the addition of a full time assistant, for approximately 50 man-days, in order to monitor a greater number of instrumented females, to recapture females for battery replacement, and to assist in searching for and capturing non-instrumented brooding hens. In addition a collateral study of predation and prey availability could be initiated to determine whether there is a correlation between the availability of small mammals and nest predation pressure.
- (9) Consider the possibility of placing poisoned sharptail or pheasant eggs in each nest to determine the specific mammalian predators involved in nest predation.



LITERATURE CITED

- Allard, H. A. 1947. Light intensity studies in Canaan Valley, West Virginia  
Castanea 12: 63-74.
- Ammann, G. A. 1957. The prairie grouse of Michigan. Michigan Dept. of Conserv.  
Tech. Bull., Lansing. 200 pp.
- \_\_\_\_\_. 1963. Status and management of sharp-tailed grouse in Michigan. J.  
Wildl. Mgmt. 27: 617-633.
- Artmann, J. 1969. Personal communication. Entomology, Fish and Wildlife  
Building, Univ. of Minnesota, St. Paul.
- Bach, R. N. and R. Stuart. 1942. North Dakota upland game nesting studies,  
1941. Fed. Aid in Wildl. Restoration Proj. 7R, North Dakota State Game  
and Fish Dept., Bismarck. 41 pp.
- \_\_\_\_\_. 1943. North Dakota upland game nesting studies, 1942. Fed Aid in  
Wildl. Restoration Proj. 7R, North Dakota State Game and Fish Dept.,  
Bismarck. 58 pp.
- Baumgartner, F. M. 1939. Studies on the distribution and habits of the sharp-  
tailed grouse in Michigan. Trans. N. Am. Wildl. Conf. 4: 485-490.
- Bent, A. C. 1963. Life histories of North American gallinaceous birds. Dover  
Publ., Inc., N.Y. 490 pp. (Reprint of Bull. 162, U.S. Natl. Museum.  
Smithsonian Inst., 1932).
- Bernhoft, L. S. 1969. Reproductive ecology of female sharp-tailed grouse  
(Pedioecetes phasianellus jamesi Lincoln) and food habits of broods in  
southwestern North Dakota. M.S. Thesis. Univ. of North Dakota, Grand  
Forks. 96 pp. (unpublished).
- Blus, L. J. and J. A. Walker. 1966. Progress report on the prairie grouse  
nesting study in the Nebraska Sandhills. Nebraska Bird Rev. 34: 23-30.
- Brander, R. B. 1968. A radio-package harness for game birds. J. Wildl. Mgmt.  
32: 630-632.
- Brown, R. L. 1963. Progress on a sharp-tailed grouse population study in two  
mixed prairie regions of northcentral and southeastern Montana. Montana  
Fish and Game Dept., Carleyle. 5 pp. (Mimeog).
- \_\_\_\_\_. 1965. Techniques developed and employed in a study of female sharp-  
tailed grouse ecology in Montana during 1963, 1964, and 1965. Paper  
presented at fourth biennial meeting of the Western States Sage Grouse  
Comm., Walden, Colo., June 22-24, 1965.
- \_\_\_\_\_. 1966a. Response of sharptail breeding populations to annual changes  
in residual grassland cover. A contrib. from Fed. Aid in Wildl. Restor-  
ation Proj. W-91-R, Montana Dept. of Fish and Game. Paper presented at  
Forty-sixth annual Conf. of Western Assoc. of State Game and Fish Comm.,  
Butte, Mont., July 12-14, 1966.

- \_\_\_\_\_. 1966b. Sharp-tailed grouse population study. Fed. Aid in Wildl. Restoration Proj. W-91-R-6 and W-91-R-7, Montana Fish and Game Dept., Helena. 23 pp. (multilith).
- \_\_\_\_\_. 1967. Sharptail grouse population study. Fed. Aid in Wildl. Restoration Proj. W-91-R-8, Montana Fish and Game Dept., Helena. 39 pp. (multilith).
- Bue, I. G., L. Blankenship, and W. A. Marshall. 1955. The relationship of grazing practices to waterfowl breeding populations and production on stockpond in western South Dakota. Trans. N. Am. Wildl. Conf. 17: 396-414.
- Bump, G. 1950. Wildlife habitat changes in the Connecticut hill game management area. Cornell Univ. Agr. Exp. Sta. Memior 289, Ithaca, N.Y. 75 pp.
- Buss, I. O. and E. S. Dziedzic. 1955. Relation of cultivation to the disappearance of the Columbian sharp-tailed grouse from southeastern Washington. Condor 57: 185-187.
- Cartwright, B. W. 1944. The "crash" decline in sharp-tailed grouse and Hungarian partridge in western Canada and the role of the predator. Trans. N. Am. Wildl. Conf. 9: 324-329.
- Cebula, J. J. 1966. Radio-telemetry as a technique used in greater prairie chicken (Tympanuchus cupido pinnatus) mobility studies. M. S. Thesis. Kansas State Univ., Manhattan. 61 pp.
- Craddock, G. W. and C. L. Forsling. 1938. The influence of climate and grazing on spring-fall sheep range in southern Idaho. U.S. Dept. Agr. Tech. Bull. 600. 43 pp.
- Darrow, R. W. 1945. Relation of buffer species abundance to fox predation on grouse nests. Trans. N. Am. Wildl. Conf. 10: 270-273.
- Dill, H. H. and W. H. Thornsberry. 1950. Cannon-projected net trap for waterfowl. J. Wildl. Mgmt. 14: 132-137.
- Doll, A. D. 1955. Sharp-tailed grouse management: progress report. Wisconsin Cons. Bull. 20 (12): 21-23.
- Dorney, R. S. 1959. Relationship of ruffed grouse to forest cover types in Wisconsin. Wisconsin Conserv. Dept. Tech. Bull. 18. 32 pp.
- Dyksterhuis, E. J. 1944. Adaptation of "A field method of judging range utilization" by M. H. Deming. U.S. Dept. Agr. Soil Conserv. Service. Lincoln, Neb.
- Edminster, F. C. 1954. American game birds of field and forest/their habits, ecology and management. Chas. Scribner's Sons, N.Y. 490 pp.
- Emlen, J. T., Jr. 1956. A method for describing and comparing avian habitats. Ibis 98: 565-576.
- Evans, K. E. 1968. Characteristics and habitat requirements of the greater prairie chicken and sharp-tailed grouse - a review of the literature. Conserv. Research Report No. 12. U.S. Dept. Agr., Forest Service. U.S. Government Printing Office, Washington, D.C. 32 pp.



- \_\_\_\_\_, and D. L. Gilbert. 1969. A method for evaluating greater prairie chicken habitat in Colorado. J. Wildl. Mgmt. 33: 643-649.
- Evans, R.M. 1961. Courtship and mating behavior of sharp-tailed grouse (Pedioecetes phasianellus jamesi Lincoln) M. S. Thesis. Univ. of Alberta, Canada. 134 pp. (unpublished).
- Frary, L. G. 1958. The 1957 prairie grouse habitat survey. Fed. Aid in Wildl. Restoration Proj. W-17-R-11, South Dakota Dept. of Game, Fish and Parks, Pierre, 6 pp.
- Grange, W. B. 1948. Wisconsin grouse problems. Wisconsin Conserv. Dept., Madison. 318 pp.
- Gross, A. O. 1930. Personal communication, p. 292. In Bent, A. C. Life histories of North American gallinaceous birds. Dover Publ., Inc., N.Y.
- Hamerstrom, F. N., Jr. 1939. A study of Wisconsin prairie chicken and sharp-tailed grouse. Wilson Bull. 51: 105-120.
- \_\_\_\_\_. and F. Hamerstrom. 1951a. Grouse of the brushlands. Wisconsin Conserv. Bull. 16(10): 7-9.
- \_\_\_\_\_. and F. Hamerstrom. 1951b. Mobility of the sharp-tailed grouse in relation to its ecology and distribution. Am. Midland Nat. 46: 174-226.
- \_\_\_\_\_. and F. Hamerstrom. 1961. Status and problems of North American grouse. Wilson Bull. 73: 284-294.
- \_\_\_\_\_. 1963. Sharptail brood habitat in Wisconsin's northern pine barrens. J. Wildl. Mgmt. 27: 739-802.
- Hanson, H. C. and W. C. Whitman. 1938. Characteristics of major grassland types in western North Dakota. Ecol. Monogr. 8: 57-114.
- Hart, C. M., O. S. Lee, and J. B. Low. 1950. The sharp-tailed grouse in Utah: its life history, status, and management. Publ. 3, Fed. Aid in Wildl. Restoration, Utah State Dept. of Fish and Game. 79 pp.
- Henderson, F. R. 1964. Grouse and grass / twin crops. South Dakota Conserv. Digest 31(1): 16-19.
- Higby, L. W. 1969. Personal communication. Wyoming Fish and Game Commission, Lander.
- Jackson, W. W. 1963. Prairie grouse movement study, 1962. Fed. Aid in Wildl. Restoration Proj. W-75-R-5, South Dakota Dept. of Game, Fish and Parks, Pierre. 10 pp (mimeog).
- Janson, R. G. 1953. Prairie grouse habitat survey, 1950-1952. Fed. Aid in Wildl. Restoration Proj. W-17-R-7, South Dakota Dept. of Game, Fish and Parks, Pierre. 9 pp. (mimeog).
- \_\_\_\_\_. 1955. Prairie grouse hunting / past, present, and future. South Dakota Conserv. Digest 22(10): 8-9.

- \_\_\_\_\_. 1956. The sharp-tailed and prairie grouse in South Dakota. Special Report, South Dakota Dept. of Game and Fish and Parks, Pierre, 30 pp. (unpublished).
- Jenkins, D. 1961. Social Behavior in the partridge (Perdix perdix) Ibis 103: 155-188.
- Johnson, C. E. 1934. Recollection of the prairie chicken and the sharp-tailed grouse in northwestern Minnesota. Wilson Bul. 46: 3-17.
- Jones, R. E. 1960. The life form concept applied to prairie chicken habitat in Oklahoma. Oklahoma Acad. Sci. Proc. 40: 134-136.
- \_\_\_\_\_. 1968. A board to measure cover used by prairie grouse. J. Wildl. Mgmt. 32: 28-31.
- Klett, A. T. 1957. Banding and marking methods in studying seasonal movements of the sharp-tailed grouse in Morton County, North Dakota. M.S. Thesis. Utah State Agr. Coll., Logan. 53 pp. (unpublished).
- Kobriger, G. D. 1964. Prairie grouse on two sandhills refuges in Nebraska. M.A. Thesis. Univ. of Missouri, Columbia. 131 pp. (unpublished).
- Kuck, T. L. 1966. Pheasant radio-transmitter study. Fed. Aid in Wildl. Restoration Proj. W-75-R-7 and 8, South Dakota Dept. of Game, Fish and Parks, Pierre. 30 pp.
- Lacher, J. R. and D. D. Lacher. 1964. A mobile cannon nettrap. J. Wildl. Mgmt. 28: 595-597.
- Lack, D. 1933. Habitat selection in birds. J. Animal Ecol. 2: 239-262.
- Leopold, A. 1933. Game Management. Chas. Scribner's Sons, N.Y. 481 pp.
- McEwen, L. C. and R. L. Brown. 1966. Acute toxicity of dieldrin and malathion to wild sharp-tailed grouse. J. Wildl. Mgmt. 30: 604-611.
- \_\_\_\_\_, D. B. Knapp, and E. A. Hilliard. 1969. Propagation of prairie grouse in captivity. J. Wildl. Mgmt. 33: 276-283.
- McKenzie, J. V. 1963. Big game browse studies. Fed. Aid in Wildl. Restoration Proj. W-67-R-2, Game and Fish Dept., Bismarck.
- MacMullan, R. A. and L. L. Eberhardt. 1953. Tolerance of incubating pheasant eggs to exposure. J. Wildl. Mgmt. 17: 322-330.
- Marshall, W. H. and M. S. Jensen. 1937. Winter and spring studies of the sharp-tailed grouse in Utah. J. Wildl. Mgmt. 1: 87-99.
- \_\_\_\_\_. 1963. Studies of movements, behavior, and activities of ruffed grouse using radio-telemetry techniques. Univ. of Minnesota Prog. Report to Nat'l. Sci. Found. 30 pp. (Multilithed).
- \_\_\_\_\_, and J. J. Kupa. 1963. Development of radio-telemetry techniques for ruffed grouse studies. Trans. N. Am. Wildl. Conf. 28: 443-456.
- Mathisen, J. 1963. Prairie grouse habitat and plans for management on the Shewen National Grassland. U.S. Forest Service, Cass Lake, Minnesota. 9 pp. (Mimeog).

- ttson, O. 1963. Wisconsin prairie chicken management, a progress report. Wisconsin Conserv. Dept., Madison. 6 pp. (Mimeog).
- Mosby, H. S. (Ed.). 1963. Evaluation of habitat, pp. 52-74. In Mosby, H.S. (Ed.). Wildlife Investigational techniques, 2nd ed. The Wildlife Society, Washington, D.C.
- Mueggler, W. F. 1962. Range resources and management problems in northern Idaho and northeastern Washington. U.S. Forest Service Intermountain Forest and Range Exp. Sta. Res. Paper 68. 31 pp.
- Newman, D. E. 1959. Sharptails: a land management problem. Wisconsin Conserv. Bull. 24(4): 10-12.
- Peterle, T. J. 1954. The sharp-tailed grouse in upper peninsula of Michigan. Ph.D. Dissertation. Univ. of Michigan, Ann Arbor. 260 pp. (unpublished).
- \_\_\_\_\_. 1956. Trapping techniques and banding returns for Michigan sharp-tailed grouse. J. Wildl. Mgmt. 20: 50-55.
- Podoll, E. 1955. Prairie grouse habitat study, 1953-1955. Fed. Aid in Wildlife Restoration Proj. W-17-R-10, South Dakota Dept. of Game, Fish, and Parks, Pierre. 12 pp.
- \_\_\_\_\_. 1957. Prairie grouse habitat survey, 1956. Fed. Aid in Wildlife Restoration Proj. W-17-R-11, South Dakota Dept. of Game, Fish, and Parks, Pierre. 5 pp.
- Folderboer, E. B. 1942. Cover requirements of the eastern ruffed grouse in northeast Iowa. Iowa Bird Life 12: 50-55.
- Robel, R. J. 1966. Behavior and mobility studies on the greater prairie chicken of northeastern Kansas. Yearbook of the American Philosophical Society, 1966: 365-368.
- \_\_\_\_\_. 1970. Possible role of behavior in regulating greater prairie chicken populations. J. Wildl. Mgmt. 34: 306-312.
- \_\_\_\_\_. J. N. Briggs, J. J. Cebula, N.J. Silvy, C. E. Viers, and P. G. Watt. 1970. Greater prairie chicken ranges, movements, and habitat usage in Kansas. J. Wildl. Mgmt. 34: 286-306.
- Sather, J. H. 1950. A light meter for cover density measurement. J. Wildl. Mgmt. 14: 138-143.
- Saunders, G. B. 1936. Game Div. Report No. 292, Michigan Dept. of Conserv., Lansing. 4 pp.
- Schladweiler, P. and T. W. Missehl. 1969. Use of mist-nets for recapturing radio-equipped blue grouse. J. Wildl. Mgmt. 33: 443-444.
- Schrader, T. A. and A. B. Erickson. 1944. Prairie chickens and sharptails. Cons. Volunteer. 7(40): 33-37.
- Schwartz, C. W. 1945. The ecology of the prairie chicken in Missouri. Univ. of Missouri Studies 20. 99 pp.

- Schwilling, M.D. 1960a. Effects of grazing on prairie chicken nesting. A contrib. from Fed. Aid in Wildl. Restoration Proj. W-33-R, Nebraska Game, Forestation and Parks Commission. Paper presented at third annual National Prairie Chicken Conf., Stevens Point, Wisconsin, Sept. 8-10, 1960, 7 pp.
- \_\_\_\_\_. 1960b. Experiments to develop livetrapping and individual bird marking techniques. Fed. Aid in Wildl. Restoration Proj. W-33-R-1, Nebraska Game, Forestation, and Parks Commission, Lincoln. 9 pp. (mimeog).
- \_\_\_\_\_. 1960c. Studies of daily and seasonal movements of prairie grouse. Fed. Aid in Wildl. Restoration Proj. W-33-R-2, Nebraska Game, Forestation, and Parks Commission, Lincoln. 8 pp. (mimeog).
- Scott, W. E. 1947. The prairie sharp-tailed grouse (Pedioecetes phasianellus campestris). Wisconsin Cons. Bull. 12(2): 26-29.
- Sharp, W. B. 1958. Microclimatic influences created by ground nesting birds. Ecology. 39: 757.
- Slivy, N. J. and R. J. Robel. 1968. Mist nets and cannon-nets compared for capturing prairie chickens on booming grounds. J. Wildl. Mgmt. 32: 175-178.
- Slade, N.A., J. J. Cebula, and R. J. Robel. 1965. Accuracy and reliability of biotelemetric instruments used in animal movement studies in prairie grasslands of Kansas. Tran. Kans. Acad. Sci. 68: 173-179.
- Smith, C. C. 1940. The effect of overgrazing and erosion upon the biota of the mixed prairie of Oklahoma. Ecology. 21: 381-397.
- Symington, D. F. and T. A. Harper. 1957. Sharp-tailed grouse in Saskatchewan. Conserv. Bull. 4, Saskatchewan Dept. National Resources. 24 pp.
- Uhlig, H. G. and W. H. Hamor. 1966. Land management for sharp-tailed grouse. In more wildlife for Recreation. U.S. Dept. Agr. Soil Conserv. Service, Lincoln. 14 pp.
- U.S. Department of Agriculture, Forest Service. 1969. Range and Environmental Handbook. In Forest Service Handbook, no date. U.S. Government Printing Office, Washington, D.C. loose leaf, n.p.
- U.S. Fish and Wildlife Service. 1955. Improvements to the cannon-net trap. Branch of Wildlife Refuges, Wildl. Mgmt. Series No. 12. 26 pp.
- Viehmeyer, G. 1941. The present status of the greater prairie chicken and sharp-tailed grouse in the sandhill region of Nebraska. Nebraska Bird Rev. 9: 1-7.
- Viers, C. E. 1967. Home range and movements of the greater prairie chicken (Tympanuchus cupido pinnatus) with notes on activities. M.S. Thesis. Kansas State Univ., Manhattan. 70 pp. (unpublished).
- Watson, A. 1964. Aggression and population regulation in red grouse. Nature 202: 506-507.
- Weaver, J. E. 1954. The North American prairie. Johnson Publ. Co., Lincoln, Nebraska. 348 pp.

West, D. R. 1959. The 1958 prairie grouse habitat survey. Fed. Aid in Wildl. Restoration Proj. W-74-R-1, South Dakota Dept. of Game, Fish and Parks, Pierre. 7 pp.

Wight, H. M. 1939. Field and laboratory technic in wildlife management. Univ. Michigan Press, Ann Arbor. 107 pp.

Yocom, C. F. 1952. Columbian sharp-tailed grouse (Pedioacetes phasianellus columbianus) in the state of Washington. Am. Midl. Nat. 48: 185-192.

Zwicker, F. 1958. North-central Washington grouse studies. Washington State Game Bull. 10: 3-4.